


Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

62.67

1227



How to Know
GOOD SEED


SCOTT'S SEED GUIDE

SCOTT'S SEED GUIDE



Points the Way
to Profitable Farming

O. M. SCOTT & SONS CO.
MARYSVILLE, OHIO



Contents

	Page
Foreword	4
HOW TO KNOW GOOD SEED—A test that should be used on every seed purchase	5
HOW TO FIGHT WEEDS	8
SCOTT'S SEED—What present users think of it	10
SOYBEANS—general description — Yellows — Browns and Blacks	12
SWEET CLOVER—History—popularity — advantages —pasturing and cutting—seed production	24
ALFALFA—general outline—feed value—kind of soil preferred—drainage—seeding—harvesting	34
GRIMM ALFALFA—Guaranteed not to winter-kill— seed now cheaper than red clover	44
THE TRUE CLOVERS—Red Clover—Mammoth— Alsike—Crimson—White Clover	47
JAPAN CLOVER—good on poor soil	56
COW PEAS—VETCH—FIELD PEAS	57-58
TIMOTHY—cheap this year	58
PASTURE AND MEADOW MIXTURES	59-60
SUDAN GRASS	63
CORN—Little Cob and other varieties	64
WHEAT—Trumbull—Fulhio—Gladden—Nigger	68
OATS—Scottish Chief—Sixty Day—Miami	69
LEGAL WEIGHT AND QUANTITY PER ACRE	70
SCOTT'S Bacteria—Friendly Workers of the Soil	71
POINTS ABOUT ORDERING	75
WE PAY THE FREIGHT	79
FREIGHT AND EXPRESS RATES	80

Foreword

BACK in 1909 we issued a little booklet called "Weeds and Seeds and How To Know Them." It told in a simple and understandable way why the selection of seed was so essential to profitable farming. Here are a few lines from that first book: "When a weed spreads rapidly over a given territory some folks like to blame birds, wind or running streams, but the start at least should be laid to the hand of man. He shuts his eyes, pays good cash for poor seed and then sows it on a splendidly prepared seed bed." Today the careful selection and thorough cleaning of seed is as essential to the profitable growing of crops as it was 18 years ago.

"Scott's Seed Guide," a sequel to "Weeds and Seeds," has never ceased harping on that vital point. Over a half million copies have spread the gospel of pure seed. In recent years this Guide has found its way into the classroom. 8000 copies were reviewed last year by students of agriculture.

We commend to you this 1927 edition of "Scott's Seed Guide" with the hope that it may be a valuable aid to you in your next 12 months of farming.

Note: Extra copies for interested friends are always available.

How to Know Good Seed

WHEN YOU buy a bushel of clover seed you expect your purchase to be practically all clover unmixed with weeds, dead grains and filth. But unfortunately a good share of the seed harvested is unfit for sowing. The great problem in seed buying is to know you are getting your supply from the best of the harvest. We urge every buyer of field seeds to use the following test in comparing different lots of seed.

TEST

Take a level teaspoonful of the seed you want to test and spread it out on a piece of white paper. Scrape to one side all the waste matter, such as dead grains, weed seeds and chaff. Compare the two piles.

One you can profitably pay for—the other means a loss, for which you pay the good seed rate.

This operation puts before the sower a simple basis of comparison between the worth of our seeds and the worth of other field seeds.

Consider that in buying a bushel of seed you get six thousand times the amount of waste matter found in a teaspoonful. Just three weed seeds in a teaspoonful means that one will be sowed on every square yard of ground when you sow at the usual rate.

To find the exact amount of pure seed, have a druggist weigh the pile of seed and then the pile of waste matter. Divide the small amount by the total weight, and the result will show the percentage of impurities in the sample.

You may be surprised to know how much worthless waste you have been buying along with your field seeds. *And remember, weeds mean a loss, even if you get them for nothing.*

50 pounds of pure, practically weedless, seed will give you a more profitable stand on a five acre field than will 60 pounds of inferior seed. That means if you paid \$15.00 per bushel for the cheap seed you could well afford to pay \$17.50 for the better quality. To the seller there is always a bigger profit on second grade seed. *To the sower there is always a bigger profit on pure seed.*

HARD GRAINS

All clover seed contains a certain percentage of hard seeds that are slow in absorbing water. This percentage is largely dependent on weather conditions at maturing time. In making tests, clover seed is left in the germinator five or six days. Of course, some of the hard grains will germinate if not removed for several days longer, and many of them, undoubtedly most of them, will grow when planted in the ground and subjected to moisture, heat and the chemical action of the soil.

While it hardly seems possible, it is a fact that seeds having the finest appearance (and containing the largest amount of plant food to give the young plants a good start) are those that ordinarily contain the greatest number of hard grains.

In buying seed it should be borne in mind that *freedom from weed seeds is of greater importance than germination*, providing, of course, that this is not unusually low.

NO WEEDS!

Allow me to congratulate you on the quality of grass seed you sell, have had perfectly wonderful success in getting a good stand when others fail, and have not found a single weed seed in any seed bought from you.

F. L. AKERS, *Ellison, W. Va.*

GERMINATION

To make a germination test, remix the seed and count out, say, 200 seeds. Be sure to take them just as they come, and do not choose the best grains, for the object is to find out what percentage of the total seed will grow. The seed may be planted in a box of moist dirt or sand, or may be put between two blotters or strips of cotton flannel, placed on a plate covered with another plate upside down. This prevents evaporation. Keep the blotters moist, but not in water, and as near the temperature of 70 degrees as possible. Examine the seeds each day and see how they are germinating. Weak or slow germination indicates that the seed will make weak plants. Sprouted seeds may be removed each day if desired. Some seeds require a longer time to germinate than others.

The proper germination periods are as follows:

Clover seed between three and six days.

Timothy and Red Top between five and eight days.

Orchard Grass between six and fourteen days.

Kentucky Bluegrass, fourteen to twenty days.

If you want to go into the matter of seed testing a little more thoroughly, send for Farmers' Bulletin No. 428, from which much valuable information can be obtained.

UNUSUAL SUCCESS WITH ALFALFA

You will remember I wrote you several weeks ago telling how good alfalfa seed purchased from you came through the awful dry spell of hot weather we had in this part of the country. We all wondered how your seed can stand such dry weather and do so good. We expected to lose the stand when weather remained so dry. My friends around here are much interested in your seed.

GEORGE W. FRIES, *Chambersburg, Pa., R. F. D. No. 3*

How to Fight Weeds

THERE isn't anything mean enough that can be said about weeds. Words can't express your and our contempt of them and of the losses they inflict. They stand in the way of many profitable harvests, adulterating the crop and preventing a thick, uniform growth.

In every Seed Guide published since 1920 we have enumerated the respects in which weeds inflict losses. Here they are again:

Weeds rob crops of moisture and valuable soil elements.

Weeds crowd plants out of space and light.

Weeds may harbor plant diseases.

Weeds are often poisonous and most of them taint milk.

Weeds may prevent the drying of hay or grain in the shock.

Weeds constitute an adulterant in harvested products.

Weeds often interfere with harvesting.

Weeds add to the expense of harvesting and take dockage from the grain.

Weeds may interfere with the system of crop rotation.

The real time to fight weeds is discussed in the section called "How to Know Good Seed." It is easier to carry on the battle described there when it is simply a case of refusing to buy inferior seed. One slip in seed buying is enough to cause weed trouble for years to come. *Prevent weeds* by sowing pure seed. That beats fighting them. But we may as well concede the existence of weeds and the following suggestions may prove helpful in combating a few common varieties.

CANADA THISTLE

CANADA THISTLE is a native of Europe. It was introduced into Canada in impure farm seeds just as it is now being sent all over this country. It is easier to eradicate during drought. Cultivation to destroy it should not be undertaken when the land is wet. All plants must be completely destroyed, as, if only a few survive, the ground is soon reinfested. Probably the best method of extermination is to starve the underground parts by destroying the top growth as it appears.

On small areas the plant may be smothered out by covering with tar paper, or boards. These may be covered with dirt or a stack of hay or straw.

Canada Thistle cannot stand shading and is often exterminated by a smother crop. Alfalfa is being used for this purpose. In addition to shading the Thistle, the continuous cutting keeps it from producing seed. The alfalfa should be seeded heavily.

WILD CARROT

Wild Carrot is a biennial, thus it can be gotten rid of if the plants are not allowed to mature seed. It grows more abundantly on hard, dry soil, on hillsides and in old meadows. If soil is badly infested a cultivated crop such as corn, potatoes or beans should be grown until the wild carrot disappears.

It causes very little trouble in fertile soil or in cultivated fields.

Wherever this weed is abundant an extra effort should be made to build up the fertility of the soil.

BUCKHORN

Buckhorn is a perennial, propagating itself chiefly by seeds.

Buckhorn can only be gotten rid of by cultivation and by sowing clean seed.

Badly infested fields should be plowed up and worked under a short rotation of crops.

Buckhorn is usually found on thin soils or on land deficient in organic matter.

Scott's Seed

We could fill the remaining pages of this Seed Guide with reasons for our firm belief that Scott's Seed will satisfy you every year better than any other seed on the market. But it isn't our desire to devote that much space to *our* opinions. More convincing are such comments as these by present customers:

"Received the seed. It is fine. I showed it to a farmer who had bought from a local dealer. He said, 'Well, I would rather pay \$5.00 per bushel more for seed like yours.'" We are fully expecting that man's order this year, for no doubt his experience with the seed has been even more disappointing than its appearance.

Consider this also: *"Here is a real compliment for you. I wrote to our Experiment Station addressing a personal friend who is on the staff. I asked where we were likely to get the best seed. In listing the names of four firms in his reply he added; 'O. M. Scott and Sons Company are the best we have found.'"* Coming from a state Experiment Station, of which we have sold thirty-eight, this testimonial is especially gratifying. If Scott's Seed satisfies those whose special training enables them to accurately test seed, surely the farmer who purchases it has little to fear.

Freedom from weeds is a respect in which we invite the most careful comparison of Scott's Seed with any other. You don't want weeds. More than that, you don't want to pay for weed seeds at pure seed prices. In commenting upon Scott's Seed, a West Virginia customer says, *"allow me to congratulate you on the quality of seed you sell. Have had perfectly wonderful success in getting a good stand with it. I have not found a single weed in any seed bought from you."*

Extract from Scott's Soybean Catechism

Why should I grow the soy bean? It is a legume, therefore a nitrogen gatherer. It produces a large yield of seed, an excellent quality of forage that is relished by all farm animals, is easily grown for harvest, is drought-resistant, and is practically free from insect enemies and plant diseases.

How should the seed bed be prepared? The same as for corn. Like corn, the soy bean readily responds to extra cultivation. Plow early and deep. Harrow in the spring to prevent growth of weeds. Disking is sufficient if the crop is to follow wheat after harvest, provided the ground is fairly mellow.

How much moisture does this crop demand? One of the best qualities of the soy is that it is drought-resistant. However, they are not sensitive to excess of moisture. Drainage is not required.

Can soys be used in rotations? Yes, they are the most satisfactory legume, especially in short rotations. They may be used as a whole season crop as in Corn, Soys, Wheat and Clover or Grass rotation; or they may be used as a part season crop following small grain as, for instance, wheat.

How can I have a legume in the ground all the time? Plant soy beans with your corn. Soys alone the next year for hay, pasture, or seed. Follow with wheat in which you sow clover.

What is the feeding value of this hay? Cut at the proper time soy bean hay contains more digestible nutrients, more carbohydrates than any other hay. Fed to dairy cows the Tennessee Exp. Station found that it produced 12% more milk and 18% more butter fat than alfalfa hay.

What is the feeding value of soy bean seeds? It contains about 40% protein and 18% fat. Needless to say its feeding value is very high.

Scott's Soybean Catechism will be sent to anyone upon request. It asks and answers 36 vital questions about the crop.

Soybeans

THEY occupy the prominent position in our Guide this year for three reasons. First, their contribution to farming justifies it. Second, they will be a foremost substitute for high priced Red Clover, sharing that position with Sweet Clover and Alfalfa. Third, Scott's Soybeans lead the field in point of quality and volume of distribution. Of the early varieties sold direct to users, we believe our sales will head the list.

We want our present customers to know and tell their friends about *Soybean Headquarters*.

Soybeans

PIPER AND MORSE, in their very interesting book, "The Soybean," say that over fifty names are used in oriental countries for the soybean. One of the names used in China is "Sou." No doubt the name "Soy" has been derived from it.

The Soybean is one of the oldest plants utilized by man. The first mention of the soybean in this country was in 1804. The Perry Expedition in 1854 brought to this country two varieties, but not until late years has the bean become well known.

The Soybean becomes more popular each season and is, in most sections, gradually taking the place of Cowpeas. Soybean seed contains both of the vitamins necessary to life.

In buying Soybeans it is well to remember that quality counts for much. Beans often contain wild morning glory and other undesirables. Many beans, too, are put on the market before they are well cured. In such cases the germination is apt to be low.

Because we handle beans in such large quantities, have a thorough knowledge of the best producing sections and are familiar with the many different kinds so as to know their best adaptations, we feel prepared to render a little more intelligent service than most firms selling Soybeans. These same advantages enable us to sell at reasonable prices. As far as quality is concerned, we believe you will not find Scott's Soybeans excelled.

Clover is so often a failure that it is necessary to find some legume that pays as large returns and at the same time is a sure crop. Soybeans not only gather more nitrogen from the air than clover, but have a larger root system. Thus even though the crop is saved for hay or is pastured instead of being plowed under, the benefit to the ground is as great as from clover, as is indicated by the large yields of wheat and corn following a well inoculated crop of Soybeans.

The Soybean is used for green manure, hay, forage, grain and silage.

It is probably the most desirable leguminous catch crop.

Invaluable to fill in with in case of failure of clover seeding or other spring crop.

The Soybean requires from 90 to 150 days to mature. Some of the varieties will mature as far north as Northern New York, but the plant having originated in a warm climate, many varieties will not ripen except in the South.

It will grow in poor soils and increases the fertility of the land by means of the nodules on its roots. It endures drought, and stands excessive moisture fairly well.

It is more frost resistant than corn or field beans. Light Spring frosts will not kill the young plants and they are seldom injured by frosts in the fall when maturing.

The plant is moderate in its fertilizer requirements. Lime, applied previous to seeding, should be used on acid soils, though liming is not as essential as with other legumes.

As might be expected, trials made at experiment stations show that for milk and butter production Soybean hay is nearly as good as alfalfa hay. Cattle will leave corn to eat it.

The forage produced by the Soybean is higher in protein than any other annual crop of equal yield.

Experiment stations are urging that Soybeans take the place of oats in the rotation. By planting corn and beans together, then Soybeans alone, either for hay or grain, followed the next year with wheat and clover, a legume is on the ground continuously. As an average crop of Soybeans, if inoculated, will store up as much as 125 pounds of nitrogen per acre, it can be seen what they will do for soil improvement when used in this way.

SEEDING

In seeding Soybeans the seed bed should be prepared about the same as for corn, cultivating at intervals in order to kill weeds, for, like alfalfa, young Soybean plants are easily crowded out by a rank growth of weeds. They should

be planted about corn planting time, but not until all danger of frost is past, and ground is warm. The seed should be planted not over 1 to 1½ inches deep. If a seed crop is expected it is best to plant as early as possible, but for hay or green manure even into August is not too late, the variety used making some difference. Although solid drilling requires more seed to the acre, and in wet seasons there may be some annoyance from weeds, many growers prefer this manner of planting for hay, silage or green manure crops. From 4 to 8 pecks are used. Laboratory experiments in New Jersey point to an increased or intensified use of atmospheric nitrogen where legumes are planted close together. When so planted there is possibly a greater recovery of nitrogen from the air per acre. Thick seeding is best for sandy soils. Planting in rows saves seed and permits cultivation.

Planted after wheat or oats are removed they will usually produce a paying hay crop. In any case they will be invaluable for soil improvement or for pasture.

In weedy ground it is best to plant in rows, but Soybeans drilled solid can be cultivated with a weeder or slant-tooth harrow. This should be done during the middle of the day when the beans are dry and tough. Do not cultivate in this way until the beans are three inches high, nor after they have reached a height of eight inches.

For seed the beans can be drilled in rows and cultivated. A grain drill can be used for drilling in rows by covering up the necessary feed holes. This requires from 15 to 25 pounds of seed according to the size of the beans. The rows should be 28 to 30 inches apart, with the beans 2 or 3 inches apart in the rows. When a corn planter is used, some recommend attaching a shoe at the planter runner for regulating the depth. For solid drilling use the oats feed of an ordinary grain drill adjusted to drill, say, 75 pounds per acre. Solid drilling for seed production is becoming increasingly popular, using in some varieties as much as two bushels per acre.

CULTIVATION

As the entire bean is pushed out of the ground in sprouting, in case a heavy crust forms, it may be necessary to harrow the ground lightly before the beans sprout, or even as they are coming through the ground, but this should be avoided if possible. Unless weeds are getting the best of the beans they should not be disturbed until three or four inches high, when they are quite tough. When planted in rows the beans can be cultivated until blossoms appear, but should not be disturbed after this.

IN CORN

When planted with the corn Soybeans may be "hogged off" or "lambd off," or cut with a binder for putting in the silo. When the former, the lambs or pigs are turned in when the corn is about ready to cut for fodder and the Soybeans have begun to ripen. Pigs up to 50 or 60 pounds and the lambs will eat the beans without injuring the corn if it is desired to harvest it. If the entire crop is to be pastured, sheep or hogs of any size may be turned in. They take on fat and improve in general condition wonderfully. It is a good plan, if convenient, to feed Soybeans to stock a few days before they are put into the field, so that they will become accustomed to, and eat the beans readily, otherwise the larger animals will favor the fresh corn. Beans are so rich in protein that it is not necessary to feed tankage or any other protein feed.

For hogging off or for silage the beans are planted right in the rows with the corn. A special attachment can be gotten for the planter that makes it possible to do the whole job at once, or the corn may be drilled first, at the regular depth, and then the drill filled with beans, going back over the corn rows, being careful to plant the beans no deeper than one inch. The corn is planted about 18 inches apart, and the beans about 6 inches apart in the rows. This requires 6 to 10 pounds of beans. If the planter has a fertilizer attachment the beans may be mixed with the fertilizer or with dust, drilling the mixture as fertilizer. Corn and beans may be mixed together and drilled, but this is not a very satisfactory arrangement.

Soybeans should be planted in all corn fields when possible to utilize them, unless the fields are inclined to be very weedy. As they gather nitrogen from the air and the corn gets some of this, the growth of the corn is not lessened, and even though it were, the benefit to the ground and the extra forage would more than offset the loss. At least one ton of silage or soiling is added to the field. Many report yields of two tons, a large profit at practically no expense.

We sell a good many Soybeans for broadcasting in corn just before the last cultivation. Under normal conditions this crop is worthwhile for hogging down or for soil improvement. Some plant them in corn each year for soil improvement, so that corn can be grown on the same field year after year. Late in the season we usually have a few varieties left that we can sell at a special price.

FOR SILAGE

The addition of the Soybeans gives silage a much greater feeding value, as they contain 145% more digestible protein and 40% more fat than the corn silage. Of course, Soybean silage contains a very much greater amount of protein when the beans are allowed to form in the pods. It should be kept in mind that, like corn, different varieties of beans have different dates of maturity, and that the variety to be selected is the one that not only makes a maximum growth of vine but also develops beans by the time the corn is ready for putting in the silo. If the beans are grown in separate fields three loads of corn run through the cutter, followed by a load of Soybeans makes a well-balanced feed. They can be used much riper than for hay, as they go into the silo without curing, and the juices of the corn soften the stems. The harvesting is done with a binder just as the corn alone would be harvested.

The addition of Soybeans to the silage makes the purchase of oilmeal and tankage unnecessary and greatly increases the flow of milk.

Cornell University in a series of experiments found that non-leguminous plants grown with legumes contained a great deal more protein than when grown alone. As an example, oats grown with field peas contained 7% more

protein than oats grown alone; timothy with red clover 44% more. It seems reasonable to suppose that corn grown with Soybeans should contain more protein than when grown by itself, especially if they are inoculated.

FOR HAY

Tall varieties are always most desirable for hay as many of the smaller kinds bear pods and leaves too close to the ground for harvesting.

The Wilson and Virginia varieties have slender stems thus making the most desirable, and perhaps the most abundant hay of any of the northern varieties.

Soybean hay in curing will stand more unfavorable weather conditions than red clover or alfalfa. The plants should be cut when the pods begin to fill and a few yellow leaves are showing. Well-matured hay may not be so palatable, but is more easily cured. When once started the harvesting should be completed in as short time as possible, as the leaves fall rapidly when ripe. They may be cut with a mower and left on the ground until wilted; then raked up and placed in tall, loose cocks for a week or ten days. A good method, however, is the use of the self-binder, setting the small bundles into cocks to cure.

FOR SEED

The seed cures to best advantage on the stalk, so beans should not be cut until absolutely necessary to prevent loss from shattering. A good guide is to wait until most of the leaves have fallen off. If the beans are well ripened it is possible to thresh in a day or two after cutting. If the beans are not allowed to get thoroughly ripe—and some varieties must be cut early or too many beans will be lost—the beans should be kept in cocks until thoroughly cured, otherwise the seed may be damaged when stored in bins or sacks. Cut when dew is on the ground. Some of the large growers of seed cut with a binder just as soon as the top leaves turn yellow, allowing the small bunches to lie three or four days, according to the weather, and then put in small shocks, reshocking at the end of ten days if the weather is favorable. In four or five weeks the beans are

thoroughly cured. This leaves a large amount of leaves on the straw, most of which are lost if the beans are allowed to ripen thoroughly. Some are now using the Southern Soybean harvesters, which gather the beans from the standing stalks.

THRESHING

An ordinary grain separator can be used by removing the regular concaves and using a blank or board. The speed must be cut down so as to avoid splitting the beans. Special bean separators can be purchased at reasonable prices. Soybean straw is relished more than any other straw by sheep, cattle and horses.

We find that a great many of our customers are using corn shredders for threshing their beans. The only change necessary is to loosen the snapping rolls and raise the back part of the shredder about four inches. Feed slowly about one load of beans per hour and keep the screens clean. While this operation is slower than using a separator, practically no beans are cracked.

INOCULATION

There is no question at all but that Soybeans should be inoculated. They may grow nearly as well without inoculation, but will do this at the expense of the soil. When inoculated the roots become filled with large nodules, which make them the ideal crop for soil building. There is also little doubt that when they are inoculated the protein content of the plant is greater. For reasons stated before, we believe it is especially important to inoculate Soybeans that are to be planted with corn. In 99 cases out of 100 "No inoculation means no nodules."

A man who has used Scott's Bacteria on Soybeans for many years recently wrote as follows: "I always use Scott's Bacteria twice on the same field. This year I had an average of $15\frac{1}{4}$ bushels per acre, at least three to five bushels better than the average in this county." (Scott's Bacteria paid its cost five times over.)

Scott's Bacteria in bushel size cans at \$1.00 is very

cheap when it comes to inoculating Clover, Alfalfa, Vetch, etc. On Soybeans the comparative cost is greater and to encourage a more general use of our cultures, we now offer a five bushel size can at a special price of \$3.50 postage paid. Since our cultures are especially effective on Soybeans, this low price should appeal to all large growers.

SOYBEANS IN MIXTURES

Soybeans may be mixed with cow peas, sorghum or sudan grass, making a balanced forage; about ten pounds of sudan grass or fifteen pounds of sorghum with three pecks of Soybeans broadcasted, make hay that is easily taken care of.

YELLOWS

MANCHU

The Manchu was introduced from Manchuria in 1913. The plants are erect and bushy and mature in 110 days. This is by far the most popular early variety and has almost completely taken the place of other early kinds such as Ito San, Early Brown and Black Eyebrow. It produces a large yield of seed and considering its earliness is an excellent hay bean.

ITO SAN

While this small yellow bean has been one of the most popular early varieties, growers have discarded it for the Manchu.

MIDWEST

This bean, formerly called the Hollybrook, is one of the best liked varieties. It gives good results when used for hay, seed or silage. It matures in about 115 days.

A K

The A K is a pretty good substitute for the Midwest and is much more plentiful.

DUNFIELD

This variety is becoming popular in Indiana where it is used in place of the Manchu. It is about five days later

and has the advantage of holding its leaves longer. It produces about the same amount of seed and hay.

MAMMOTH YELLOW

This is a late southern grown variety, maturing in about 145 days. It will not ripen beans north of the Ohio River, but is grown to some extent for hay as far north as the New England States. We recommend an earlier kind for either hay or silage.

BROWNS

VIRGINIA

This bean was selected from another kind in 1907. The tips are twining—probably more so than any other variety. It grows quite tall with slender stems, making it one of the most desirable beans for hay and for planting with the corn for silage. The vining tendrils cling to the corn. It is particularly desirable on poor ground where height is desired but on the richest soil it is sometimes inclined to lodge. It matures in about 125 days.

BLACKS

Most of these have fine stems and leaves, and are desirable for hay and for silage, but many growers prefer the larger and coarser yellow beans for the latter purpose. In blacks the same variety often goes under several different names. We have found the Wilson, Sable, Sooty, Jet, Peking and Arlington to be very much alike. Experiment Station reports vary greatly as to time of maturity and other characteristics.

PEKING OR SABLE

The Peking, often called Sable, was brought from Peking, China, in 1907. While not so well known as some of the other black varieties this is a desirable bean and will go a long ways in planting.

WILSON

The Wilson is one of the best known and one of the most popular all around medium early varieties. Introduced from Manchuria in 1906. After having been grown on the government farms for two years it was sent out to experiment stations in 1909. In 1911, a special selection was made by W. J. Morse, soybean specialist of the Department of Agriculture. This selection was so desirable that it was soon in great demand.

The plant is slender and erect which makes it just the bean for hay and silage. As it matures in about 120 days it is early enough to make hay in almost any part of the country.

The Wilson is the best all around variety for hay and silage. The stems are fine and it produces a larger quantity of hay, of better quality than the Manchu and other yellow varieties. In buying any black beans it is well to be on the lookout for wild morning glory. We use extreme care to remove all of this weed from our beans. The quality this year is above the average.

EBONY

This is a stout, erect growing, bushy variety with fine stems. It is somewhat earlier than the Wilson and produces a less amount of hay. Matures in 115 to 120 days.

BLACK EYEBROW

Another early variety maturing in 105 to 110 days. Adapted for hogging off and for early hay and seed production.

OTHER VARIETIES

While this year we are not listing the Early Brown, Ohio 9035 and Medium Green we may be able to supply such of our customers as want these or any other variety.

MIXED

Usually we can furnish mixed beans at a lower price than separate varieties.

FIRST AND SECOND CHOICE

We attempt to furnish practically all varieties of Soybeans and will have a few not listed here but inasmuch as so many beans are much alike, we suggest, especially later in the season, that in ordering, first and second choice be given.

We have the most improved machinery for cleaning them and believe we are furnishing better beans than can be purchased most places.

We give dates of maturity for Ohio. The number of days required for the ripening of beans will vary somewhat with the locality and weather conditions.

Ask for "Soybean Catechism."

LAREDO

A black bean that has become popular during the last few years. As it is only a little earlier than the Mammoth Yellow it is not adapted to northern sections.

"I harvested twenty-five bushels of Soybeans per acre from the poorest field on my farm." This statement was made in our office by a grower from whom we purchased three hundred bushels of beans. As a result of this first experience he assured us he would increase his acreage next year and that many neighbors would follow his example.

Sweet Clover

COMING into more general use every season, the seed is not nearly as plentiful as last year. For this reason and because everyone seems to think it will be used largely in place of Red Clover, the price is somewhat higher. The quality is very good.

This is an ideal year to get started with Sweet Clover. Those who have already used it recognize its value and sow it in increasing quantities. If you have never sowed Sweet Clover we suggest reading the following pages carefully.

Biennial Sweet Clover

The Universal Plant

SWEET CLOVER is a universal plant, because it can be grown in nearly all parts of the world. It is raised in every State in the United States. In many clover-producing sections that have suffered from Red Clover diseases, it is replacing this legume. Sweet Clover seems almost entirely free from diseases.

There are a large number of varieties of Sweet Clover, but only three are of value in this country, namely, White Biennial (*Melilotus alba*), Yellow Biennial (*Melilotus officinalis*), and Annual White Sweet.

Grundy County is a smaller variety of White Biennial Sweet Clover.

While the white, on account of its larger growth, is the most popular variety, the yellow biennial is preferred by many on account of its finer and more branching stems. It seems to be a larger producer of seed. It has the advantage of maturing about two weeks earlier than the white.

When Sweet Clover is mentioned without any special variety being named, it is always understood that the biennial white is meant.

Its extensive root system enables it to gather the little plant food remaining in worn-out soils. When inoculated, the abundant root nodules store nitrogen and the decaying roots add humus. The deep penetration of the longer roots improves the drainage and after a crop of Sweet Clover the soil is always more friable and mellow, so that following it a good crop, even of corn, may be raised, though no profitable crop could be grown before.

Observe the uncultivated spots where Sweet Clover appears to grow the best. Organic matter, or humus, seemingly is not necessary, but these spots always contain lime and the ground is always hard. This indicates the method that should be employed in order to be sure of a good stand

of this valuable plant. A firm seed bed should be prepared and lime must be applied if the soil is acid.

Inoculation is almost necessary.

Freezing weather does not injure bacteria. The sun's rays do not prevent inoculation when seed is sown on top of the ground as enough bacteria will stick to the under side.

Scott's Bacteria is guaranteed for Winter as well as Summer use.

Sweet Clover prevents erosion on hillsides. It practically never freezes out during winter or spring. It is a weed-killing crop because of its rank growth. It builds up worn-out pastures and meadows. It will carry several times as much stock as ordinary pasture land. It contains more protein than Red Clover. Unlike alfalfa, it is not injured by pasturing. Plowing it up is easy, for although the roots are quite large, they are soft and decay rapidly.

Bee keepers for years have recognized the value of Sweet Clover. The honey from Sweet Clover is of good color and flavor.

As a universal plant it leads even alfalfa, for it will grow not only in any climate, but on soils where alfalfa fails. Where it is at first impossible to get a stand of alfalfa, Sweet Clover will furnish a profitable yield and at the same time prepare the ground so that there will be no difficulty in establishing alfalfa permanently.

For four or five weeks after germination, Sweet Clover makes but a slow growth above the ground, but the root system is developing, getting ready to meet unfavorable conditions, and as is the habit with biennials storing up food not only for this but also for the next season's growth.

The tap roots during the first season will sometimes reach a depth of thirty inches. The top growth is rapid after the plant has once become established.

FOR GREEN MANURE

According to recent experiments conducted by the Illinois Experiment Station, Sweet Clover offers the most promising indications that the problem of supplying a cheap source of nitrogen can be solved.

Sweet Clover possesses the following characteristics, which make it far superior to other crops as a green manure for nitrogen production :

1. It is adapted to almost any climate.
2. Hardiness to cold or drought.
3. Resistance to disease and weeds.
4. Large production per acre at the proper time for plowing under.
5. Rapid decomposition.
6. Deep rooting habit makes impervious subsoils porous.
7. Obtains plant food from insoluble minerals more readily than other crops.

A crop that grows rapidly in early spring, and decomposes rapidly, makes an ideal green manure. It also keeps in the soil large amounts of plant food by its rapid growth. Besides, Sweet Clover, when properly inoculated, gathers nitrogen from the air. It stores in its roots great quantities of nitrogen, which are available as soon as growth starts the second year. When turned under, the leaves decay rapidly; the roots and stems more slowly. The three different parts represent three sources of nitrogen to be drawn on over a considerable length of time.

Few soils are able to meet large crop demands for nitrogen. Since Sweet Clover will supply nitrogen at an insignificant cost, it could well be utilized by most farmers as a green manure crop.

One soil may contain twice as much nitrogen as another and still produce less crops, because the nitrogen is resistant to decay. A small amount of active nitrogen in a poor soil may produce more than a larger amount of inactive nitrogen in a rich soil. It is most important to keep as much nitrogen as possible in the active state of decomposition when it is needed by crops. If this is done by inoculated legumes, then the amount of nitrogen that should normally, or always, be in the soil, is at the same time preserved.

Sweet Clover's ability to do this should not be overlooked. To conserve nitrogen may be as important as its addition.

Investigations by Messrs. Whiting and Richmond showed that one ton of water-free, or about six tons of green Sweet Clover (roots and all), furnish as much nitrogen as twenty tons of farm manure.

The tops on a water-free basis averaged 4.38 per cent nitrogen, which means 87.6 pounds of nitrogen per ton of dry weight. The roots weigh as much as the tops and the nitrogen content is about the same, which indicates more than 170 pounds of nitrogen.

Sweet Clover, when used as green manure, will add to, conserve, and make available the nitrogen of the soil.

PASTURE

It is claimed by some that an acre of Sweet Clover will furnish pasture for five or six times as many animals as will the ordinary mixed grasses. While some stock will refuse Sweet Clover at first, they all soon get the habit, especially if turned into the field when the plant is small and tender.

During the first year grazing can start when the plant is about six inches high and can be continued until late in the summer.

When seeded on wheat fields during the winter or early spring an abundance of fall pasture is available when most fields are affording very little feed.

The second year Sweet Clover makes a quick, early growth and may be pastured sooner than any other plant. If a seed or hay crop is wanted it can be pastured until the middle of June, for grazing really benefits the stand by causing the plants to stool and make a larger number of branches.

When a field is used for pasture alone enough animals should be grazed to keep it eaten reasonably close. Then there will be a constant supply of small, tender shoots. Should the plants become coarse the pasture can be clipped to stimulate the growth of fresh shoots; setting the mower knife eight inches high.

Do not be afraid of pasturing too closely. Usually the more stock you turn on Sweet Clover the better.

If stock are removed about two months before heavy frost the pasture will reseed itself.

Besides furnishing the earliest pasture, it thrives during the hot, dry summer months and makes some growth after the first frosts.

One acre will furnish pasture for at least twenty shoats.

There is less danger of bloat than from alfalfa pasture.

As stock crave a dry feed when pasturing on Sweet Clover, it is often desirable to have a stack of straw or hay in the meadow.

The white is best for pasture as it stays green longer.

CUTTING FIRST YEAR'S GROWTH

The first year the crop should be cut for hay at about the time growth ceases. There are two important reasons for cutting at this time. In the first place, this season's growth does not become woody and it is safe to wait until the maximum growth is made. Second and more important, there is no danger of injuring the plants by cutting too close, because the first growth of the second year starts from the crown. Shortly before the end of the growing season the crown buds are noticeable, after which it is safe to cut. This point is not generally well understood.

The fact that buds for new growth are all ready to start at the beginning of spring, along with its vigorous root system, makes it produce pasture so much earlier than other plants.

The first year, do not clip Sweet Clover during the summer unless absolutely necessary.

If the plant is clipped to kill ragweed or other weeds, the cutter bar should be set quite high, for after being clipped there is no further growth from the main shoot, but dependence must be had upon the lateral branches for pasturage or for hay. As many of these should be left below the point of cutting as possible.

CUTTING SECOND YEAR'S GROWTH

"While the first crop in the second year comes from the crown buds, the new branches which produce the second crop of the second year come from the buds formed in the axils of the leaves on the lower portions of the stalks which constitute the first crop. These branches usually commence growth when the plants are about twenty-four inches high. In fields where the stand is heavy and where

the lower portions of the plant are densely shaded, these shoots are soon killed from lack of necessary light. The branches which are first to appear and which are first to be killed are those closest to the ground. It is therefore very important when cutting the crop to cut the plants high enough from the ground to leave on the stubble a sufficient number of buds and young branches to produce a second crop.

"In fact, the stand should be cut several inches above the young shoots or buds, as the stubble may die back from one to three inches if the plants are cut during damp or rainy weather."

We quote from U. S. D. A. Farmers' Bulletin No. 820. This explains very clearly why care should be used in cutting Sweet Clover. We advise reading of this bulletin as well as Mr. Coe's other Sweet Clover Bulletins, No. 797 and No. 836.

Hay cut the first year is fine-stemmed and palatable.

Hay must be made the second year before the bloom buds appear, as the plants become woody about this time.

The hay is cured in the same manner as Red Clover or Alfalfa, but being more succulent a longer time is required.

As in handling other clovers, the idea is to get rid of the water gradually instead of allowing the leaves and stems to be burnt by the sun. This saves the leaves, the most valuable part of the plant. The hay should lie in the swath until well withered and then be raked into windrows. The next day, if sufficiently dry, it must be put in cocks and cured. The cocks should be of such size that they can be loaded in one forkful in order that as few leaves as possible be lost.

SEEDING

From a labor-saving standpoint, at least, probably the best time to sow Sweet Clover is during the winter any time from January to April on corn ground or other bare ground. Freezing and thawing will bury the seed and cause the hard grains to germinate, there being quite a large percentage of these in any Sweet Clover that has not been scarified. Zero weather will not kill bacteria.

Even when the laboratory test shows the germination

of Sweet Clover to be quite high it will not respond to moisture as quickly as Red Clover, more rain being needed even when the seed has been scarified. For this reason growers are sometimes surprised to find that they have a poor stand of Sweet Clover, while in a neighboring field sowed with Red Clover at the same time, the growth has been entirely satisfactory.

We quote a letter received from the Department of Agriculture: "I have yours of the 7th and in regard to the failure of Sweet Clover to germinate or to live after germination, when the seed bed is not firm this appears to be entirely a matter of moisture supply. A loose seed bed is dry especially when a period of protracted dry weather follows sowing. A firm seed bed is in contact with the moist subsoil. Just why Sweet Clover should be so much more sensitive than other clovers we cannot say. There are peculiarities of all varieties that we simply know exist, but we are not in a position to explain them."

A lack of moisture just at the time of germination or immediately after the young sprout is appearing seems to affect Sweet Clover more seriously than other clover. However, there is a very small percentage of failures in the seeding of Sweet Clover.

Remember that Sweet Clover seed should be only lightly covered.

A firm seed bed is important, as it is in contact with the moist subsoil, so where necessary to plow, if possible the ground should be plowed in the fall and harrowed down, the seed being broadcasted during the winter months. However, seeding may be done in April or May on a well prepared, firm seed bed with just enough loose soil to cover the seed. Like any other clover, Sweet Clover may be seeded on wheat or rye in the spring, or with oats or barley.

August seeding is not desirable in most sections, for then the plant lasts but one growing season and does not reach its largest development. Many sow during June or July; some as late as the last of August. The Virginia Experiment Station recommends sowing in August for pasture and hay crops the following year.

While Sweet Clover, once it is established, is very

drought-resistant, the plants when young must have an abundance of moisture on account of the deep-growing roots.

Too heavy seeding means that young, small branches and leaves on the lower part of the stem will be killed by the crowding. This necessitates higher cutting and makes less and coarser hay.

Inasmuch as Sweet Clover has a hard seed coat, scarifying has proved to be a profitable operation.

It requires ten or fifteen pounds of this seed to the acre; when the unhulled seed is used fifteen or twenty pounds. The scarified seed is best for summer or fall sowing, for the germination is more rapid. For winter sowing possibly the unscarified, hulled seed is to be preferred. We furnish this or the scarified seed at the same price.

SEED PRODUCTION

When a seed crop is to be saved, Sweet Clover should be first pastured or cut for hay. The stock should be removed from the pasture early enough to assure the maturing of seed; if hay is made, the cutting should be reasonably early, thus assuring a larger number of branches. Of course the mower should be set high, as new shoots will not come from the crown, but from the axils of the lower leaves.

The mower, however, is not as satisfactory for cutting Sweet Clover as the binder, for the reason that too much handling is necessary. The self-rake reaper is best, but a binder can be equipped at small expense for handling the clover economically and with small loss of seed. Corn harvesters are sometimes used if the growth becomes too large to be cut with the binder.

A very thin stand produces a surprising quantity of seed when neither clipped nor pastured.

Cutting should be done when about three-fourths of the pods have turned dark, and only when damp from dew or rain, as the seeds shatter easily. For this reason, when cut with a mower, the swath should not be run over.

When cut for hay or pastured, the late plants are smaller and are harvested more easily, for the binder can be used. This facilitates handling and makes it possible to

use a huller, otherwise it is sometimes necessary first to thresh the coarse straw and then run the seed through the huller.

IN MIXTURE

A mixture about like the following is popular: Sweet Clover, four pounds; Alfalfa, four pounds; Alsike, two pounds; Timothy, four pounds.

RENOVATING OLD PASTURES

Worn-out pastures are often successfully renovated by disking in the fall and sowing a few pounds of Sweet Clover during the winter. Not only is the amount of pasturage increased by the Sweet Clover, but the grasses will be improved owing to the addition of humus and nitrogen furnished by the Sweet Clover. The same plan may be followed in the spring, but not as successfully. The Sweet Clover should be drilled in.

HUBAM

While this annual legume has been abused and its value to the farmer questioned by many authorities, we believe that it will take a permanent place in agriculture.

It fits in well for an emergency or regular hay and pasture crop, and can be sowed on wheat or oats.

GRUNDY COUNTY SWEET CLOVER

Grundy County Sweet Clover was developed in Grundy County, Illinois. It is a biennial but does not grow nearly as tall as either the White or Yellow biennial. The stems are finer and it has become quite popular in many sections. Ripens about three weeks earlier than the common White Sweet Clover.

PRICES HIGHER, BUT QUALITY COUNTS

Your prices are very much higher than the Louisville, Kentucky market, but I haven't been having very good results with their seed.

C. A. BRIDGES, *Campbellsville, Ky.*

Alfalfa

ITS popularity has never waned. We have never heard a serious criticism of Alfalfa. It provides an abundance of rich hay and greatly benefits the ground while doing it. Scott's Alfalfa Seed will stand any quality test. The following pages contain helpful authoritative information for old as well as new growers of this money-making legume.

Alfalfa

Compared with other clovers, Alfalfa is low in price this season and the quality is excellent. We call particular attention to the fact that Grimm Alfalfa is cheaper this year than Red Clover.

A FIELD of alfalfa should last for four or five years before being crowded out by Blue Grass or other native grasses. This being a fact, it is surely wise to take extra pains to learn just what kind of alfalfa seed is being sown. It is easy to buy imported alfalfa. There is lots of it, and the price is lower. Argentine seed, at least, has an attractive appearance, for it is usually free from weed seeds, plump and bright. Unfortunately Experiment Stations have found that it winter-kills, the same as other imported alfalfa. It in no way compares with home grown alfalfa in hardiness.

If you are not sure about the seed that you are offered, why not send a sample to the Department of Agriculture. They can readily identify it for you.

Even in purchasing American alfalfa it is well to know in what states the seed was produced.

ALFALFA'S WORST ENEMY

It is a fact accepted by all Experiment Stations that weeds are the thing to consider when sowing alfalfa. The seed bed should be clean, and the seed itself free from weed seeds. If weeds are abundant, the young alfalfa plants will be smothered out. To get a good stand of alfalfa, two things are obviously necessary: land free from weed seeds, and alfalfa seed free from weed seeds. The first year weeds are alfalfa's worst enemy. Aggressive weeds can ruin a good stand.

In sowing alfalfa, it is a question not so much of pounds per acre as of quality of seed and proper soil conditions. Seven pounds of seed that is pure and of strong vitality is better than fifteen pounds that contain weed seed.

The expense of preparing land is the same no matter what kind of seed is sown. The actual difference in the cost per acre between the best and the poorest seed is very

little, but the cost of care and of harvesting is greatly increased if weedy seed is sown. It is most unwise to sow weeds with one hand and fight them with the other.

In the first part of this book we have demonstrated many of the losses due to weeds and have shown that inferior seeds thoughtlessly sown may mean the spreading of weeds all over your own and neighboring farms.

FEED VALUE

Protein is the element which to a large extent determines the amount of beef or milk a given feed will produce. The market value of any feed is based on the amount of protein it contains.

Nearly all stock feeds fall short of the proportion necessary to proper balance, but alfalfa furnishes this needed element in abundance.

You should not buy protein if you can produce it on your own farm.

Too often on farms where alfalfa is not raised, an unbalanced ration is fed owing to the high cost of concentrated feeds. A large part of the otherwise necessary cash outlay for high protein feeds can be avoided by the use of alfalfa, the most economical ration balancer at the command of dairyman or stockman.

Alfalfa and corn make not only the cheapest but the best balanced ration. To feed corn and other grains alone wastes starch, and animals do not thrive as well as when along with the corn a rich, palatable feed, such as alfalfa, is fed.

Circular No. 25, Michigan Experiment Station, says: "A ton of alfalfa hay contains 46.5 pounds of nitrogen, 12.2 pounds of phosphoric acid and 35.8 pounds of potash, and that its total present value, if purchased as commercial fertilizer, would be about \$10.80."

Alfalfa hay can be raised for much less than \$10.80 per ton, in fact it can be sometimes purchased for this price. When this is the case, one is actually getting the hay for almost nothing, because in returning the manure to the fields, only 25.3% of nitrogen, 22.5% of the phosphorus and 12.2% of the potassium is lost.

The sowing of Scott's Seeds will keep the manure pile free from weeds.

PREPARATION OF SEED BED

It is not possible to plow timothy or blue grass sod for immediate planting of alfalfa, without having the growth of alfalfa greatly interfered with by the grass. A cultivated crop, such as corn, should be raised the year before; then it is unnecessary to plow the ground unless it is foul or of a very heavy character.

If plowing is thought best it should be done in the fall, especially if early spring seeding is contemplated, for in seeding alfalfa the loose, open seed bed, such as is prepared in plowing a short time before seeding, should be avoided. It takes nearly six weeks for plowed ground to settle for alfalfa seeding. Otherwise, capillarity, or the power of the soil to draw water to the surface is interfered with.

The surface should be cultivated until the soil is as fine as a well-prepared garden. This top covering prevents evaporation, and thus keeps the soil warm, besides leaving the seed in complete contact with the soil, which makes plant food easily available when the seed sprouts.

The young plants are likely to die in poorly prepared spots, these spots eventually becoming weed distributing stations. Hollows become filled with water and ice, which may kill the alfalfa.

COMPETITION OF WEEDS. POOR SEED

The ground always contains weed seeds, and for this reason, thorough cultivation at intervals to kill them as they sprout is advisable. After going to this trouble, it is surely unwise to sow seed that contains weed seeds which will sprout at the same time as the alfalfa, and probably overcome it.

LACK OF FERTILITY

While alfalfa is a deep feeder, drawing its food from greater depths than most plants, it is more tender than other clovers when young, and needs encouragement. Well rotted manure is the best fertilizer, but usually not available. Fresh manure, on account of weed seeds, should be applied to the preceding crop, or before plowing, the weeds

being destroyed by frequent cultivation up to seeding time.

FERTILIZERS

If it is impossible to use manure, commercial fertilizers should be used, especially on poor soils. We like bone meal best. Any fertilizer used should contain a large amount of phosphorus. Alfalfa gathers its own nitrogen, if inoculated.

ACID LAND. NEED OF LIME

The percentage of lime in the ash of alfalfa is almost 35, nearly twice the percentage shown by red clover, and more than seven times that in timothy. Lime is beneficial to most plants, and to the legumes it is absolutely necessary. More lime is needed for alfalfa than for red clover.

If sorrel, dock, red top or blackberry bushes thrive in your fields and clover does not, undoubtedly lime is needed. If muriatic acid poured on the soil fails to make bubbles, lime should be used. If limestone pebbles or shells are present lime need not be added. The absence of these, usually, but not always, indicates the absence of lime. Valleys are not so apt to require lime as hills.

When in doubt about lime use it, or consult your experiment station.

A large crop of alfalfa cannot be expected unless the bacteria which find their home in the nodules on the roots are present. These bacteria gather free nitrogen from the air, and pass it on to the alfalfa, but they have no use for an acid soil.

Of the three forms of lime, the one that is the most economical in your section should be used.

One hundred pounds of raw lime rock when burned is reduced to 56 lbs. of burnt lime or quick-lime. When this is water-slaked it takes up 18 lbs. of water, making 74 lbs. of hydrated lime. Therefore, 56 lbs. of burnt lime, or 74 lbs. of hydrated lime is equivalent to 100 lbs. of ground limestone.

Probably not less than two tons of ground limestone per acre should be applied, and more will not hurt. Apply as long before sowing as possible, even a year in advance.

DRAINAGE

All plants require air in contact with the roots. If there is too much water in the soil, the air is reduced and root de-

velopment retarded. A lesser top growth follows. Drainage takes off surplus water and admits air, causing circulation. Almost any wet soil, if properly drained, will raise alfalfa. The tile should be put in as deep as may be practicable.

INOCULATION

A great many scientists have been giving years of study to the subject of inoculation. As far as we know, every one of them agrees that it is necessary to introduce the proper bacteria into the soil if the best stand of alfalfa is expected. The cost of time and material is small, so it scarcely pays to run the risk of partial failure in order to save the trouble. Nitrogen is the most expensive fertilizing element. If the bacteria are supplied, alfalfa fills the soil with it at no cost.

WINTER-KILLING

Alfalfa seldom winter-kills on land well drained. Any clover is apt to winter-kill if there is little humus in the soil. Ground containing plenty of humus is porous and ventilated; there are air spaces which favor the rapid carrying-off of excess moisture. Stiff clay soils that are devoid of humus become filled with water, which in freezing weather forms ice and causes the soil to expand and heave, pulling out and breaking off the roots. Fields that carry red clover through the winter will do the same for alfalfa. If each year it is becoming harder to keep a stand of clover, put humus in the soil. Soils that contain humus are warmer than soils that do not.

SEEDING

Alfalfa may be seeded at any time from early spring until late summer.

Where winters are particularly severe, spring seeding is favored. In milder climates, due to the early growth of weeds, fall seeding is desirable.

Sowing in the corn is a good gamble, if care is used to have the field free from weeds and in good order. A one-horse drill may be used, or the seed broadcasted, followed by a small harrow. If one cultivation is omitted, the alfalfa has a better chance on account of the earlier sowing.

Alfalfa does better than Red Clover when sown in wheat fields or with oats in the spring because it will get along with less moisture. Sixty Day Oats, owing to the smaller growth, is the best variety for a nurse crop. Not over a bushel per acre should be used.

Some sow in June, but we have found that foxtail and other weeds are still likely at that time to be a serious menace, and because we don't like weeds, we prefer to sow from July 20th to August 10th. This allows plenty of time for getting the ground in fine shape and for killing the weeds by cultivation. At this time the young plants receive full benefit of moisture, sunlight and plant food, as they sprout. Late seeding has many advantages on most soils. For one thing, the land can be used for another crop before being prepared for alfalfa. The second year's production of hay will almost invariably be greater from late summer sowing in spite of the longer time for growing allowed by seeding in the previous spring.

Fertile ground does not require as much seed as poor ground. Poor seed will not go as far as good seed. Broadcasting requires more seed than drilling. Thus the amount required per acre varies, but as a general rule twelve pounds per acre will be found about the right amount. After sowing harrow lightly to cover seed and smooth out any furrows left by the drill, as a heavy rain may bury the young plants. Alfalfa seed should never be covered to a greater depth than one inch. The top soil should be left loose.

IMPROVING A POOR STAND

It is not always possible to thicken a stand where the plants are thin throughout the field, but if they are not over a year old the following plan may be used with some success: Disk or otherwise cultivate the field, and harrow after sowing. The proper time to do this is early in the spring, before the old plants start. A thin stand from spring sowing may be improved by re-sowing in the fall if the rainfall is abundant. Ordinarily, alfalfa, more than one year old, makes a growth that will shade and smother out young plants. In improving a stand of this kind, some recommend frequent clippings until the new growth gets started.

HARVESTING

Alfalfa leaves contain twice as much protein as the stems, about twenty-eight per cent in the bud stage. This should be kept in mind when the hay is being made, and every means used to save the leaves. Where the leaves shatter in feeding they should be saved. When scalded and mixed with other feed for hogs they make an economical source of protein.

The Kansas Experiment Station found that the protein content of alfalfa when one-tenth in bloom is 18.5%, when in full bloom, 14.4%. For this reason the mature hay is better for horses. But for other stock the green hay, with the larger protein content, is the more profitable. Providing new shoots have started, the best time to cut is just as the field begins to show blooms. There should be an abundance of shoots about 1½ inches long. If too long, they will be clipped by the mower, and growing time will be lost, for alfalfa grows from the end of the shoot, like a fern. Diminished yield results if the crop is cut too soon.

The basal sprouts are the best guide to cutting, but in dry seasons the new shoots are sometimes slow in appearing. In this case the blooms must be taken as the sole guide, for the feeding value diminishes rapidly as leaves are shed to protect the plant from the drought.

In wet seasons the second growth may be six or eight inches high before the blossoms appear. It is better to observe both the sprouts and blossoms before mowing. Crowding alfalfa with too frequent cuttings will weaken the roots for the reason that they will not receive enough food from the stems and leaves; and weeds and grass are not smothered out as effectually, for they will have more air and sunshine. Nothing is gained in the end by cutting a fourth crop.

If the hay is allowed to dry too rapidly this loss of leaves happens during the curing process. A little experience, however, soon demonstrates that alfalfa is easily cured.

Alfalfa may be put in the stack or mow damper than is generally supposed, that is, while the stems are still quite tough or flexible.

Alfalfa should be let lie until the leaves are wilted, then raked into windrows and then into cocks, until cured. A heavy crop may be tedded. Alfalfa may be stacked when so dry that moisture cannot be wrung out by twisting the hay.

Usually the greatest growth is from the first crop, which is nearly always harvested during a rainy time, and requires more care in handling. Later crops, harvested in mid-summer, may be cut in the morning, raked into windrows in the afternoon, and put into the mow the next day.

Any kind of hay should be exposed to the hot sun as little as possible while curing. Too long exposure bleaches the leaves of alfalfa, and causes them to become brittle and fall off. Moreover, if the leaves have been burnt in the sun, they will not absorb the water in the stems, and the hay will cure slowly and unevenly. Curing through the action of air and wind is best, therefore, alfalfa should be cured in cocks, instead of in the swath.

The greatest quantity of hay is harvested during the third or fourth season. After this weeds, grass, etc., weaken the stand, and the yield decreases.

Do not clip alfalfa in order to kill weeds until it is in blossom, or at least until the weeds are about ready to seed. This will kill the weeds in young fields. Early mowing may kill the young plants. The idea is to cut as many of the weeds and as little of the alfalfa as possible.

The alfalfa should not be allowed to form seed, for this requires plant food which should go to the roots.

PASTURING

As a general rule it is best not to pasture alfalfa under any condition, although some authorities say that it is safe to pasture when the crop is ready for making hay by turning enough stock on the field to harvest it quickly.

LEAVES TURNING YELLOW

The first growth of alfalfa sometimes shows yellow. This may be due to lack of either humus, of drainage, or of lime. Sometimes it is because the young plant is feeding in the surface soil and the condition disappears when the tap root has pushed farther down.

SOUTH DAKOTA ALFALFA

Seed from South Dakota is credited with being hardier than that grown in other states. States farther south produce seed more abundantly. The more or less limited production and the good reputation of South Dakota seed cause it to command a higher price. Usually we have experienced difficulty in procuring Dakota seed that is as free from weeds as we like to have it. However, by purchasing direct from the grower we are able this year to supply seed that will please the most particular and most experienced alfalfa enthusiast.

UTAH ALFALFA

Because in Utah conditions are just right for the development of a seed crop, production in this state is usually large. The seed moves at a lower price than that from other northern states. No doubt the seed produced in the northern part of this state is nearly as hardy as the hardiest ordinary alfalfa. There seems to be little winter-killing of fields from seed raised here, or, for that matter, from Kansas seed.

CANADIAN VARIEGATED ALFALFA

This Canada grown variety promises to become quite popular. Canadian growers claim that it is much like the Grimm. This has caused many to offer it for sale as genuine Grimm at a price far above its actual market value. The seed is usually plump and free from weed seed.

Grimm Alfalfa

We Guarantee Our GRIMM NOT to Winter-Kill

Grimm Alfalfa is now cheaper than Red Clover. It is a good time to get a field of Grimm started.

WE GUARANTEED it for the purpose of dissipating whatever doubt there may be in the minds of prospective customers concerning seed purchased in a section distant from the place where first grown. This guarantee is designed to give sowers the necessary confidence not only in our GRIMM, but in the GRIMM strain as a safe and profitable crop.

To guarantee the seed to grow after it is in the ground would be impossible, as too many things can happen. But if weather conditions permit the growth to come along all right, we will replace the seed, willingly and cheerfully, if the plants do not go through the winter. GRIMM ALFALFA should, of course, be planted in ground that will carry red clover through the winter; ground where the drainage and other conditions are not unfavorable. North of the Ohio River, alfalfa should be planted not later than August 10th because of the danger from dry weather, delaying germination so long that the plant will not get a good start before winter. We cannot guarantee GRIMM that has not reached a growth of 6 to 8 inches before it becomes dormant.

GRIMM will outstand ordinary alfalfa in wet ground, but it cannot be successfully grown on ground poorly drained. All investigators agree that it will do better on hard-pan soil than ordinary alfalfa, and it is the safest seed to sow in any part of the country.

As the Department of Agriculture and the Minnesota Experiment Station indicate, it is the inherent characteristics of the strain rather than the locality which first reared it that make GRIMM ALFALFA HARDY.

SCOTT'S SEED GOES FARTHER

The proof of genuineness is hardiness. There is no other positive proof. We GUARANTEE the hardiness of the GRIMM ALFALFA we furnish.

To guarantee GRIMM to be true to name, and to guarantee GRIMM not to winter-kill, are not at all analogous propositions. You could never prove that it was not GRIMM, but you certainly could prove it winter-killed if that were the case.

Different winter conditions cause alfalfa to kill. A rainy fall prevents the plants from becoming dormant early, making them much more likely to winter-kill than following a dry autumn.

Alternate freezing and thawing in clay or in humus-poor soil will break off the roots. Sheet ice often kills alfalfa.

GRIMM has the characters that enable it to withstand these conditions better than any other variety, and at the same time, it is thought to be more drought-resistant than ordinary alfalfa. For this reason many believe that on an average GRIMM will produce probably more hay than other strains.

Wendelin Grimm, a native of Germany, brought with him to Carver County, Minnesota, a small quantity of alfalfa seed. The seed produced exceptionally hardy plants, and when it was eventually recognized as a superior strain it became known as GRIMM ALFALFA.

Just why GRIMM ALFALFA is so much hardier than other alfalfa has never been satisfactorily settled, but the principal reason generally accepted is its low set crown which affords protection to the tenderest part of the plant. It is also aided by the branching tendency of the roots. It is, however, difficult to distinguish GRIMM from ordinary alfalfa by examining the root system. It does not show so large a percentage of branching roots as one would be led to believe from illustrations of selected plants and from some advertisements. Generally, about 40% to 70% of GRIMM roots are more or less branched, but a considerable per cent of the roots of common alfalfa also show that tendency. There is not much difference in the blossoms,

except that GRIMM when in full bloom shows a higher percentage of mixed or variegated flowers. The seeds of both are exactly alike.

GRIMM ALFALFA has a definite market value like any other standard seed. GRIMM at a very low price could not be true to name, and so would be an unwise buy.

In seed producing sections GRIMM fields are grown for seed almost exclusively, owing to the extra price which the seed brings. The weeds in an alfalfa field tend to increase more rapidly when the stand is allowed to remain for seed each year than when the field is mown regularly for hay. For this reason one should be especially careful in purchasing GRIMM ALFALFA.

SCOTT'S BACTERIA INCREASED YIELD

Three years ago I sowed one package of your bacteria on a five-acre piece of wheat that we seeded and the first hay crop was simply immense. We had at least 15 or 16 tons of clover and timothy mixed.

(Signed)

J. H. HIBLER, *Hubbard, Ohio.*

SCOTT'S SOYBEANS GAVE HIGHEST TEST

Last year I put on a variety test using your Midwest beans with eight other varieties furnished by the Department of Agriculture, and your Midwests made the highest test of the nine varieties.

A. G. HUFFMAN, *Creamery, W. Va.*

220 BASKETS PER ACRE!

The Little Cob Seed Corn I got from you went 220 baskets of corn to the acre, and shells more corn to the basket than any corn I know of that will mature here. I surely am pleased with it.

R. J. GLIEM, *No. 1, Wooster, Ohio.*

.....The True Clovers.....

ALWAYS popular but for two years seed has been very scarce. There is very little native Red, but read about the adaptability of French seed. We offer it as a necessary expedient. Native Alsike is available. It always pays to buy Clover Seed from reliable sources. You can rely on Scott's Clover. Let us do your weed-worrying for you.

Red Clover

For the second time in our experience, practically no good Red Clover seed was produced in this country except in the far west. Oregon and Idaho had some very fine looking Red Clover and a good deal of it, but all of the Idaho seed contains a sprinkling of Sweet Clover and Alfalfa. While this is not much of a detriment, it lowers the actual purity of the seed. Clover in other sections is of poor color and not as plump as it should be.

While we have never offered French Clover it will be necessary to do so this year. French Clover is stained one per cent green. Other adaptable Red Clover from other countries is stained the same way, except Canadian which is stained violet. Red Clover that is not adapted to our conditions is stained ten per cent red. As a rule the French seed has a high purity and is entirely free from Sweet Clover and Alfalfa. There is still a lot of French Clover from last year in the country that was received before the new staining law went into effect. We know that some of it is being offered as home grown seed owing to the fact that the price asked is below the market value of domestic seed.

RED CLOVER was first cultivated in Persia. It was carried to Spain and Italy in about the sixteenth century and was soon introduced into Holland. From Holland it was taken to England, and in about 1770 to Pennsylvania.

Red Clover is often called June or Medium Clover, the latter term to distinguish it from Mammoth,

Of all clovers Red is the most popular. Many feel that there is little need for other clovers where the Red can be grown. It is widely distributed through Europe and the United States and Canada, but unfortunately many soils that formerly grew Red Clover are now finding its cultivation difficult.

This is caused by one or more of the following conditions: plant diseases or insects; the exhaustion of potash; acidity; lack of nitrogen gathering bacteria; lack of humus or organic matter.

Humus provides ventilation, binds the soil together and increases its water-holding capacity, prevents winter-killing and furnishes food for bacteria.

Much can be done to make "clover-sick" soil again productive by restoring organic matter through the application of stable manure. Liberal application of lime will sweeten it and make conditions favorable for bacteria. Fertilizer rich in phosphorus and containing some potash is best for most soils. Even though Red Clover has been raised on the field in former years, in many instances it will be found advisable to inoculate.

It should be borne in mind that imported red clover is more susceptible to disease than home grown seed.

After clover is several weeks old the roots shorten and draw the crown of the plant into the soil. This protects the plant and prevents winter-killing.

While Red Clover is recognized as a biennial, if favorably located it may last four or five years. It is observed that, when sowed with Timothy, the third year from seeding the stand is about half and half; the fourth year mostly Timothy.

Red Clover has always been popular for pasturage be-

cause it starts new growth quickly when eaten off. It gives a large yield of hay which is easily cured. The second crop can be utilized for hay, pasture or seed.

Red Clover is not adapted to stiff clay or worn-out soils. On these soils Mammoth is much to be preferred.

Best results are usually obtained from spring sowing, and for best germination the seed should be drilled in. In some sections of the country where spring sowing fails, summer seeding has proved advisable. Especially is this true south of the Ohio river. The plants escape the hot, dry weather of midsummer which often kills or weakens them.

It is often the practice to cut Red Clover too late. If the bloom begins to ripen the plant is injured. If cut when just in bloom the second crop will be heavier, there will be no danger of harming the plants and the hay will be more palatable. Red Clover, after being cut, will not stand as much rain as either Alfalfa or Soy Beans. Thus it pays to cure it and get it in the mow or stack as quickly as possible. If cut in the afternoon when it contains less moisture it can be tedded the next morning and windrowed, shocked and mowed the same day. Partly cured Red Clover hay will not stand a thorough soaking.

Red Clover matures about the same time as some of the worst weeds, as for instance Buckhorn, Wild Carrot, Sorrel and Dodder. For this reason it is more difficult to find Red Clover free from weeds than any other variety.

We give especial attention to Red Clover seed because we sell more of it than any other. We have always been able to supply seed practically free from weed seeds and waste matter and we expect to continue to do it.

The difficulty of procuring good looking home grown seed and the ease of procuring good looking foreign seed

WHY IS IT NECESSARY TO SELL HOME GROWN RED CLOVER AT FOUR OR FIVE DOLLARS PER BUSHEL ABOVE THE TOLEDO MARKET?

There really are no market quotations on home grown Red Clover seed because imported seed is delivered in the transactions made on the Toledo Board of Trade.

The trader selling one hundred bags of clover seed to be delivered immediately or at some future date can purchase imported seed at several dollars per bushel less than home grown seed with the result that imported seed is used in making trades.

Just as long as home grown seed is as scarce as it has been during the past few years, and is badly wanted by the sower, it will sell far above Toledo's quoted price.

FOURTH ORDER—MORE COMING

This is my fourth order this year and will need some Alfalfa seed later. Your seeds have proven very satisfactory. I have the best alfalfa field in our township and a clover field that is a feast for the eyes. As secretary of our township farm bureau I shall endeavor to have our members obtain the best seeds possible.

J. C. CONGER, *Peninsula, Ohio.*

A few forkfuls more of hay will pay the difference between impure seed and Scott's.

It wouldn't be so bad to sow weeds if they didn't grow.

Mammoth Clover

For the second time we are compelled to withdraw our guarantee on Mammoth Clover. With the exception of one other year we have always guaranteed our Mammoth to be absolutely true to name. We think best to withhold the warrant this year because there was so little Mammoth Clover raised in the country.

MAMMOTH CLOVER, also called English, Sapling, and Pea Vine Clover, like red, is a biennial, but where soil and climate are particularly favorable, or where prevented from producing seed, it is likely to show a perennial tendency.

As Mammoth Clover makes hay about three weeks later than Red, it is better for sowing with Timothy or Red Top, Red Clover being overripe at the proper time for harvesting either Timothy or Red Top.

Mammoth grows to a greater height than Red, has larger roots that penetrate to a greater depth, and for this reason will often do well on soils where medium clover will make an unsatisfactory growth, the vigorous growth of the Mammoth enables it to gather more plant food from impoverished soils.

On poor soils Mammoth makes more desirable hay than on good because the growth is not so rank. On sandy soil it is superior to Red, which it excels as a green manure crop on account of its large growth of roots and stems. The long roots enable it to withstand drought and winter-killing better than Red.

While the hay is coarser than Red Clover hay it has the advantage of ripening later when there is less danger from rain.

If a seed crop is to be made the clover should be pastured until about June first or clipped, otherwise the plant is likely to exhaust itself in the production of stems and leaves. If the weather is especially dry, care must be used in pasturing as the plants may not receive enough growth

to produce a large seed crop. On very poor soils it may not be necessary to pasture at all.

Mammoth makes a much surer crop of seed than Red. It matures seed about three weeks earlier.

The very heavy growth usually smothers out most of the weeds and as a result we can always furnish Mammoth that is free from weed seeds. It seldom contains blasted grains.

Mammoth is supposed to be less subject to diseases than Red Clover. This may be because of its more vigorous growth and long roots which draw plant food from a great depth.

The seeds of Mammoth and Red Clover are so nearly alike that they cannot be distinguished. This likeness has resulted in much annoyance for the grower. We used to receive dozens of letters each year asking how we knew our Mammoth was true to name, most of the writers stating that they had more than once sowed Mammoth only to reap a crop of Red.

This seemed to be the common experience all over the country. We finally got tired of assuring people that they would find our seed true to name and have for several years answered this question by guaranteeing the genuineness of any Mammoth Clover purchased from us.

Alsike

Like Red and Mammoth, Alsike is a much shorter crop than last. It is also much higher in price. Inasmuch as it goes at least twice as far as Red, and some think three times as far, it is an excellent substitute. We usually have Alsike containing some White Clover that we can sell at less than the price for pure Alsike. It contains only a small quantity of the White.

Alsike seems to have been first cultivated near the village of Sike or Alsike, Sweden, and to have been introduced into England in 1834. It is not known when it was brought to America. On account of its appearance and

habit of growth it was once thought to be a hybrid between white and red, but is now supposed to be a distinct species.

While not strictly a perennial it usually remains in the ground for several years. Enough of the heads escape the mower and the grazing of stock to do much toward reseeding. It is not nearly as particular about acid soils as Red Clover and will withstand winter-killing much better. It should be used in mixtures on any type of soil where the seeding is to remain more than three years.

It is particularly adapted to wet soils, sometimes doing well in standing water.

It is free from the diseases that affect Red Clover.

The hay is finer than Red Clover and is preferred by stock but less is produced per acre.

Alsike gathers nitrogen from the air the same as Red Clover, and would be as valuable in the rotation as a soil builder except for its smaller root and stem growth.

Alsike and Timothy ripen together and the Alsike does not crowd the Timothy as badly as Red Clover does. For these reasons Alsike is preferred for growing along with Timothy. Because the Alsike does not crowd, it is often sown with Red Clover. It interferes but little with the growth of the Red Clover and should the latter fail to grow or be killed the Alsike will quite likely take its place. Often Alsike, on account of its spreading roots, will keep the Red Clover from "heaving" out. Much less Alsike than Red Clover should be used.

Except where grown for seed it is usually best to sow some other seed with Alsike, such as Timothy, Orchard Grass, Blue Grass, or Red Clover.

A good hay mixture is three parts Timothy, two parts Red, and one part Alsike.

South of the Ohio river Alsike, Red Top, and Orchard Grass make a desirable mixture for a semi-permanent pasture.

There are approximately 700,000 Alsike seeds to a pound, and 250,000 in a pound of Red Clover, so it takes much less of Alsike to sow an acre of ground.

As the seed is so small it should be lightly covered.

Probably because of acid soil, Canada Thistle, Sorrel

and Buckhorn infest many of the sections where Alsike is raised for seed, so it is well to look out for these weeds when testing samples. They cannot be entirely removed in cleaning, as many of the weed seeds will be the same size as the Alsike. This is especially true of Canada Thistle.

Owing to its smaller size Alsike is hard to clean, but, by using care in selection, we are always able to furnish seed that is practically weedless.

CRIMSON CLOVER

Crimson Clover is said to be a native of Southern Europe. It was introduced into Chester County, Pennsylvania, in 1820, but until 1880 its distribution was quite limited.

Crimson Clover is a winter annual, that is, being sown in late summer it goes through the winter in a green state, and matures its seed and dies in the spring. It will seldom stand the winters north of the 40th parallel.

Practically all of the Crimson Clover sowed in this country is imported from Europe. This means that the seed usually contains the seed of noxious weeds so that a careful examination should be made before buying. Use our test.

WHITE CLOVER

White Clover is usually called White Dutch to distinguish it from White Sweet Clover.

Many alsike fields contain White Clover. When the seed is harvested the two cannot be separated. Sometimes we have this mixed seed at prices lower than when the two seeds are bought separately.

CUSTOMERS LIKE SCOTT'S BACTERIA

My customers are all pleased with results. Your Bacteria should sell big next year.

JOHN C. WARREN, *Rose Hill, Ill.*

Japan Clover

A Blessing to the man with Poor Soil and Steep Hillsides

FOR years Japan Clover has been grown in the South and has proven itself a valuable plant as far north as the Ohio River. Many say it will do well in central Ohio, Indiana and Pennsylvania.

It can be sowed in mixtures (two or three pounds per acre) but its most common use is on thin old pastures where it should be sowed in January, February or March at the rate of five pounds per acre.

Unlike other clovers it does well on sour soil. Japan stores nitrogen in the ground so that the blue grass will re-establish itself. It will grow on any type of soil from sandy to stiff clay. On poor hillsides, washed places and in open woodlands it soon establishes itself. It affords pasture in July and August when the grasses are not growing. All animals relish it.

Japan Clover starts growing about the middle of May. As it is an annual and spreads by reseeding itself, it is adapted only to such latitudes as will furnish a season long enough to mature seed. It makes a growth of six to eight inches so can hardly be harvested for hay.

The seed comes unhulled only and ordinarily contains considerable waste matter. Scott's Japan Clover is different from that obtainable most places because it is very much cleaner.

COWPEAS

Cowpeas are being gradually replaced by soybeans and are so high in price this year as compared to soybeans, that we will probably not offer them at all.

VETCH

Two kinds of Vetch are of agricultural importance in this country: Common Vetch (*Vicia Sativa*) and Hairy Vetch (*Vicia Villosa*). While there are both Winter and Spring strains of common vetch, the Spring is the only one sowed to any extent in this country, except along the Pacific Coast. It is an annual.

Hairy Vetch, also known as Sand and Russian Vetch, is a Winter Annual. It is especially well adapted to sandy soils and will do much to build up any poor soil. It will withstand very cold weather, and is without doubt the safest and best fall sowed cover crop for northern states. It is more drought resistant than most legumes. Being a great nitrogen gatherer, it should be inoculated.

Hairy Vetch is sown alone or with wheat or rye as a supporting crop. Where the winters are severe the nurse crop is indispensable.

Grains of Hairy Vetch are hard and require more moisture to cause germination than wheat or rye. For this reason, when planted in a dry season, the nurse crop sometimes makes an excellent stand while the Vetch shows only a few plants.

Many sow Hairy Vetch in July, but the usual practice is to sow it during the latter part of August or up to the middle of September with rye. Ten to thirty-five pounds of Vetch are sown with three pecks to one bushel of rye. The more Hairy Vetch the more the soil will be improved.

If Hairy Vetch is seeded earlier than it is safe to sow rye, a better stand is assured. The rye can be planted later.

Hairy Vetch grows rather slowly in the fall, but recovers quickly in the spring, and makes an abundant growth to be plowed under, pastured or cut for hay.

Hairy Vetch may be seeded in the spring for pasture, either by itself or with a nurse crop, such as oats or barley. Sometimes it is mixed with Canada peas and oats as is Spring Vetch. In any case it will furnish excellent summer pasture.

CANADA FIELD PEAS

Field Peas are usually spoken of as Canada Field Peas, the name having been given when the plant was comparatively unknown and the seed mainly imported from Canada. However, few varieties originated in that country.

Being a legume they are soil improvers and furnish a ration rich in protein. They can be sown for soiling and fodder, and are unsurpassed for green manure. They are usually sown with oats, about one bushel of each, thoroughly mixed. This combination makes a very desirable hay or soiling crop, the yield being quite large.

Unlike cow peas they should be sown as early as possible in the spring, and do best farther north than Central Ohio.

One bushel of Field Peas, one bushel of Oats, four pounds of dwarf Essex Rape and eight pounds of Sweet Clover make excellent hog pasture that can be sown in the spring, the pigs being turned in when the oats and peas are about eight inches high. The clover may be omitted.

TIMOTHY

Timothy is one variety that is very cheap this year. There was a large crop. The quality is good, but on account of rains it is a little darker than usual.

This grass was first brought to this country from England by Timothy Hanson of Maryland in 1720.

Timothy is distinctly a grass for hay rather than pasture, as it does not take kindly to trampling and close grazing. It is our hardiest and best known grass and is a part of all mixtures.

The facts concerning seeding, harvesting, etc., are so well known that it is unnecessary to enumerate them.

Often Timothy seed contains a considerable amount of sorrel owing to the fact that both grow on acid soil. It is well to be on the lookout for this as well as for Canada thistle which is not easy to identify in Timothy seed.

In Timothy seed you will nearly always find a small amount of alsike, and quite often grasshopper specks. It is not possible to entirely remove either of these, and while

they hurt the looks of the seed, they make no difference in the quality, and should not be confused with black plantain, which is somewhat triangular and flat.

One peck is the amount usually sown per acre, or if clover is to be sown in the spring—a bushel to six acres. A satisfactory mixture is 7 pounds Timothy, 7 pounds Red, and 3 pounds Alsike.

PASTURE MIXTURES

A mixture gives a longer period for grazing, furnishes a greater variety, yields a crop richer in protein and makes a better balanced ration than would the grasses composing the mixture if sown separately. But it does not pay to sow in a mixture any grass that will not do well alone. In choosing the grasses to go into the mixture such varieties should be selected that the good qualities of one will balance points in which the other is deficient. For example, the grass that forms roots on the surface is not desirable from the standpoint of fertility; another may send its roots fairly deep, but not be as suitable for pasture grass as the other. The two make a combination well adapted to grazing and maintaining the fertility. Pasture Mixture grasses should be selected with respect to their periods of growth so that grazing may be done through the longest possible period.

A small amount of various clovers should be included in the Pasture Mixture as legumes not only feed the grasses by pumping plant food from great depths to the surface, but also supply them with nitrogen drawn from the air, and, no doubt, greatly increase the protein content of the grasses. A small amount of Alfalfa will do much towards getting the soil inoculated. White clover will grow where nothing else will and Alsike does well in wet places. Due consideration must be given to the fact that the kinds of grasses that should be used depend upon the locality. Even in a single field parts will be found that are adapted to grasses that will not thrive in the rest of the field.

Salting patches of such weeds as quack and wire grass to induce close grazing will often rid the field of these pests.

Those weeds most distasteful to cattle thrive best in meadows.

Sow pasture mixtures at the rate of 20 to 25 pounds per acre.

MEADOW MIXTURES

Meadow Mixtures, in contrast with pasture mixtures, should contain only grasses that mature at about the same date.

For reasons already stated it is more profitable to sow a mixture of several grasses, including clovers, for hay rather than to sow one kind alone, for then the roots fully occupy the ground to a considerable depth, each variety getting its food from a different level, the legumes acting as feeders for the grasses.

EARLY PASTURE AND SOILING MIXTURE

Annual pasture mixtures have become popular. More energy value from the same acreage can be procured when these are made use of as soiling crops. Canada Field Peas and Oats probably take first rank. Rye and Vetch as well as Japan Millet and Dwarf Essex Rape are also used. Any of these can be added as desired to the following mixtures which we recommend:

No. 1. 8 pecks oats, 4 pecks field peas.

No. 2. 4 pecks oats, 3 pecks barley, 3 pecks rye.

No. 3. 6 pecks oats, 5 pounds sweet clover, 5 pounds alsike clover, 6 pounds timothy.

KENTUCKY BLUE GRASS (*Poa Pratensis*)

This variety of grass is native both to Europe and to North America, and, along with two or three other similar species, is the greatest American pasture grass. Authorities are of the opinion that it is grown more or less in every state in the Union. It makes the best sod of any of our grasses and does fairly well on a wide range of soils, although it is better adapted to clay than to sandy loam. It is a very nutritious pasture grass, but has little value for hay. The fact that it is both an early spring and a late fall grower makes it valuable for grazing at both ends of the season. Kentucky Blue Grass constitutes a part of practically every lawn and pasture mixture.

Blue Grass is much cheaper this year than it has been for a long time, probably cheaper than it will be for a num-

ber of years. With Blue Grass and Timothy both low in price, one can afford to build up worn out pastures and meadows.

ORCHARD GRASS (*Dactylis Glomerata*)

Orchard Grass, known as Cocksfoot in England, is a native of Europe. Its American name is due to the fact that it is successfully grown in partially shaded places.

Orchard Grass will stand more drought than Kentucky Blue Grass, but is not especially adapted to dry land conditions. It starts very early in the spring and grows rapidly so that it is valuable in a pasture mixture. Orchard Grass is inclined to grow in tufts or bunches so that it will not permit an even sod. Although of high nutritive value, it is not relished by stock as well as Blue Grass and Red Top. It thrives best on rich, well-drained loams and makes a good growth in shady places. Twenty-eight pounds is the amount usually sown per acre.

It is seldom possible to get Orchard Grass that does not contain a considerable amount of dock and sorrel and quite often buckhorn, all undesirable weeds.

RED TOP (*Agrostis Alba*)

Red Top belongs to a class of grasses that are very widely distributed over the globe. It is a perennial which ranges in growth from a few inches to three or four feet according to the condition of soil and climate. It starts later in the spring than Kentucky Blue Grass, grows slower and matures later. Red Top is valuable for pasture and hay, but does not equal Timothy. While adapted to a great variety of soils it does especially well on wet bottoms and should always be included in mixtures for such land.

About 15 pounds of clean Red Top should be sown per acre.

MEADOW FESCUE OR ENGLISH BLUE GRASS

This deep-rooting strongly perennial grass is not better known because sowers have clung to Blue Grass, Timothy and Red Clover in mixtures. We recommend it strongly for all mixtures. Its deep roots will prevent other grasses from winter-killing.

MISCELLANEOUS GRASSES

Besides the grasses already mentioned, we are able to offer such varieties as are in general demand: Canada Blue Grass, Rye Grass, and all imported fancy grasses.

LAWN SEED

We are entering upon an age of good lawns. In the past anything that was green seemed to answer the purpose, but now we look at the matter differently.

Even Real Estate firms have come to realize that an attractive lawn helps to sell a house. It isn't that the grass in itself is so valuable, but a humble looking house with a big velvety lawn becomes a *home* and homes are worth more than houses.

To make and maintain a beautiful lawn is not the easiest thing in the world, but our booklet, "The Seeding and Care of Lawns," tells how to get the most out of your efforts and expense. We will gladly send you a copy on request.

It is "good lawn insurance" to sow Scott's Lawn Seed. We do not believe there is a purer or better mixture on the market. Considering that fact you can't make a more economical purchase. The price is fifty cents per pound, postage paid. A pound will go twice as far as a pound of ordinary lawn seed.

MILLET

The term Millet takes in a large group of forage plants, the Foxtail being the one most widely known in this country. To the Foxtail group belong the Common, Hungarian, Golden, and Japanese varieties.

GOLDEN MILLET

This is the most largely used variety, being grown to a great extent in the West, most of the seed coming from that section.

Thirty-five to fifty pounds per acre.

HUNGARIAN MILLET

This is smaller and earlier maturing than the Golden. The hay is somewhat more desirable as it is not as coarse. Thirty-five to fifty pounds per acre.

DWARF ESSEX RAPE

This valuable plant has been extensively grown in this country during recent years only.

The cost of sowing is very small as only four or five pounds per acre are required.

It grows from 1½ to 4 feet high and makes a large amount of forage for sheep, hogs or cattle.

SUDAN GRASS

Sudan grass has become increasingly popular because it will grow on almost any kind of ground, will withstand drought, and yields abundantly. It will make a crop on less moisture than any other grass. It is an annual of the sorghum family, but produces several times as much feed as Sorghum or Millet. It is fully equal to Timothy as a feed. It is used for pasture, soiling, ensilage and hay. While it is a very heavy grower, the hay is not coarse and it is relished by horses, cattle and hogs.

When all of the smaller grasses are withered on account of dry weather, Sudan furnishes plenty of green, rich pasture. It is particularly desirable in any ground which does not hold the ordinary pasture grasses. It should be sown about two weeks after corn planting time, with a grain drill or broadcasted. Fifteen to twenty pounds per acre are required unless drilled in rows to be cultivated for seed, when four pounds per acre are sufficient.

The first crop is ready for hay in from sixty to eighty days. It should be cut when in bloom, although cutting may be done a little earlier when the plant has started to head out. The second crop should be ready in forty-five days. Two crops are all that can be expected in the North although as many as four are harvested in the South. The plants stool remarkably.

INOCULATION WORKED FINE

Enclosed find check covering the amount due you for the soybean inoculation. It worked fine.

DON R. LAWRENCE, *Leonardsburg, Ohio, R. F. D. No. 1.*

Corn

OLD-FASHIONED farmers believed that stock did better on yellow corn than on white. Experiment stations, basing their conclusion on chemical analyses, told the farmers that there could not possibly be any difference; for stock feeding yellow corn and white corn were of precisely the same value.

The Wisconsin Agricultural Experiment Station has made scientific tests recently, using the effect of the two on the growth of pigs as a basis of judgment. These tests have invariably shown that under fall conditions yellow corn supplemented with whey, skimmed milk, or linseed meal, is the more profitable, and that pigs in the "dry lot" fed on white corn with skim milk made less growth and often became rickety, (a condition sometimes called paralysis or rheumatism). Pigs were carried from 50 or 60 pounds to 200 pounds weight on yellow corn with a greater gain per day and at a less cost than shown for a similar growth on white corn.

Fat-soluble vitamins that are necessary to the growth of all young animals are found in yellow corn but not in white. These vitamins are abundant in butter fat, are found to a limited extent in tankage and meat meal, but seem to be lacking in whey, skimmed milk, and linseed meal. Animals on pasture and green hay do not suffer from vitamin deficiency; the trouble is entirely with pigs under winter conditions where they cannot obtain green feed. Tankage and meat meal used as supplements to white corn do not entirely make up the deficiency.

The Wisconsin experiments showed that white corn and skim milk could be fed to pigs by adding a small quantity of chopped legume hay. They used alfalfa chopped in a silage cutter with an alfalfa screen. A small amount of hay is suggested because the digestive tract of young pigs is not adapted to dry roughage.

Five per cent of alfalfa added to yellow corn and tankage showed a gain in rate of increase and a reduction in cost. The further addition of linseed meal to this ration was still more satisfactory.

In closing their discussion they recommended that "every hog and pig in Wisconsin get legume hay each day when not on pasture, unless they are fed yellow corn and dairy by-products."

White corn seems to be as good as yellow for horses, dairy cattle, beef cattle and sheep, if they are fed ordinary well-balanced rations, including plenty of good green-colored hay or good silage. * * * *

The best insurance against loss of your labor in growing corn is being certain that you plant good seed. No one ever produced a good corn crop from poor seed.

A maximum yield can be expected only when there are no weak or missing stalks. To plant a bad ear means about 900 weak, barren or missing stalks to the acre.

Cultivation, fertility of soil and drainage affect the production of corn but the crop depends first upon the selection of seed.

It is necessary to keep down such rank growing weeds as foxtail, cockle, ragweed, etc., as well as the smaller weeds in order to save the moisture and fertility of the soil for the corn, which needs both in abundance.

Experiments have shown that nearly always shallow cultivation is preferable to deep. In no case should corn that has reached a height of 2 or 3 feet be cultivated deeper than 4 inches. By the time the corn has reached this height the roots have spread from hill to hill. Probably 2 or 3 inches is deep enough to kill the weeds and at the same time miss the roots.

Cultivation aerates the soil and keeps it warm by preventing the evaporation of moisture.

The butt grains are fertilized first, and the pollination of the grain proceeds in regular order towards the tip. Owing to this delay in development the tip grains are thought to vary from type more often than the grains on the rest of the ear.

Experiments conducted by the Kansas Experiment Station in the field show that 90% of the middle grains produce plants while only 86% of the butt grains and 70% of the tip grains produce plants.

Thus corn should always be carefully butted and tipped

by hand before grading, for no grader will entirely eliminate these undesirable grains. If they are not removed the planter is likely to drop the seed unevenly which will cause a smaller stand, as a uniform number of grains to the hill or space must be planted to insure the largest yield.

Corn should ripen early enough to escape frost and late enough to make use of all favorable growing weather.

The smaller early types are now believed to be more favorable for filling the silo than the larger ensilage corns. To give best results silage must have a larger percentage of nutritive value than is found in the immature sappy fodder of southern sorts. The ears should be ripe enough to be well dented and not too soft.

A few years ago we grew and began to sell a not very well known, and unnamed variety of corn that had been grown in a near-by county for a great many years. This corn gave better satisfaction in different parts of the country than any we had ever handled. Its popularity increased as it received a larger distribution. Owing to the characteristic small cob we named it Little Cob and are selling large quantities each year.

LITTLE COB YELLOW DENT

This is an early corn with a very small cob. The ears average 9 inches in length. Although a yellow corn, every cob is white. The grower who began raising it forty years ago thought he liked white cobs better than red. It is carefully selected, high-bred corn and as we sometimes find a single ear of Flint in a field we suppose there was a mixture of Flint in the original selection. This may account for its very small cob and the fact that it matures early and thoroughly.

For a quick-ripening, all-round corn we know of none that will give better satisfaction. We recommend it for early ensilage as the stalk contains more leaves than other varieties. If you are not entirely satisfied with the corn you are growing, we suggest that you give this variety a trial. It matures in about 110 days. We have about dis-

continued selling late varieties. The percentage of failures is too great. We believe there is no variety that will surpass Little Cob as a general all-round corn.

EARLY YELLOW DENT

This is a medium size early corn and quite popular in many sections of the country.

EARLY CLARAGE

This yellow corn matures in about 110 days and never fails to ripen. The ears are fair size. It is a very satisfactory yielder.

IMPROVED CLARAGE

A name carried by many different types of yellow corn. Ours matures in about 115 days.

McGINNIS

The McGinnis is an early white cap variety maturing in about 110 days.

ENSILAGE CORN

In ensilage corn we can supply the Virginia or Blue Ridge type. This is one of the best ensilage corns. It makes a large growth of thick succulent fodder.

We can also furnish the Eureka.

LITTLE COB IS POPULAR

One of our customers has made this report: "Quite a few in this section tried your Little Cob Corn this year. The quality and yield were good and all are well pleased."

EARLY WHITE CORN

We have an excellent early variety of White Corn. While we do not sell as much white corn as we formerly did, still some like it, especially for poor ground.

Wheat

OWING to the fact that a threshing machine goes from one farm to another always carrying at least a few grains from each place, it has been extremely difficult to supply wheat that is unmixed, but by specializing in the excellent varieties developed by the Ohio Experiment Station, we are able to furnish seed that is practically pure. The three varieties, Trumbull, Fulhio and Gladden have become very popular in other states than Ohio owing to the fact that they are all good milling wheats and large yielders.

Wheat should be sown two bushels to the acre, as it has been shown through many tests that where this amount is used more profit is realized than where six, seven or nine pecks are sown. There is absolutely nothing in the claim that a peck or half-bushel of certain varieties is enough for an acre.

TRUMBULL

This beardless variety is a selection from Fultz. It is a smooth, semi-hard wheat, much liked by millers; and what is more interesting to the grower, a large producer. Probably more of it is sold in Ohio and neighboring states than any other kind.

FULHIO

This is another Fultz selection with a stiff, easily handled straw, and is a high yielder.

GLADDEN

This is a pure line selection of Gypsy, developed by the Ohio Experiment Station. It has a very stiff straw, is a large yielder, and as a milling wheat is above the average. It is bearded, with a white, smooth chaff and red kernel.

NIGGER

This is a bearded, hard wheat very much favored by millers. It is a large yielder and usually tests higher per bushel than any variety we know of. The grains are larger and plumper than most other winter wheat. We cannot recommend this variety too highly.

Oats

IN THIS section we produced a very satisfactory crop of oats this year. The weight is good but in some varieties the color is not as bright as usual owing to wet weather.

Oats are usually sown two bushels to the acre.

SCOTTISH CHIEF

For a number of years we have sold these oats and as they are produced plentifully in this county, we can sell them at a very satisfactory price. They come to our elevator in large quantity and we select the best crops for seed. They originated from a car of Canadian oats which we distributed ten years ago. The oats are heavy and plump, and large producers.

SIXTY-DAY

Not only is this the earliest Oats, but one of the largest yielders. Owing to its extreme earliness it will make a crop where later varieties will fail. The straw is short and does not lodge, which makes it particularly desirable for a nurse crop. It has great feed value for the reason that the hull is thin and light. One of the best varieties for a soiling crop used in connection with field peas.

MIAMI

This is what the Ohio Experiment Station says: "Originally known as Ohio No. 6203. Originated as a pure line selection from the Siberian variety in 1906. Spikelets 2-3 grained, kernels white, awns weak, straw medium stiff. Average height, 40 inches; tillering power good; medium maturity. Has the highest ten-year average yield of any variety on Experiment Station farm at Wooster."

FULGHUM

This is one of the best known and highest yielding early varieties.

BARLEY


The best known bearded Barley is the Oderbrucker. We also have beardless Barley of good quality.

Legal Weight and Quantity Per Acre

KIND OF SEED	Pounds Sown		Weight
	Per Acre		Per Bu.
Timothy	10 to	15	45 lbs.
Alfalfa (broadcast)	10 to	15	60 lbs.
Alsike	5 to	8	60 lbs.
Red Clover	10 to	15	60 lbs.
Crimson Clover	10 to	14	60 lbs.
Sweet Clover (hulled)	12 to	15	60 lbs.
Sweet Clover (unhulled)	15 to	20	30 lbs.
Red Top Fancy (solid seed)	14 to	20	14 lbs.
Blue Grass	30 to	35	14 lbs.
Orchard Grass	25 to	30	14 lbs.
Meadow Fescue	20 to	24	24 lbs.
Tall Oat Grass	20 to	30	14 lbs.
Lawn Grass Seed	60 to	80	
Canada Field Peas (with oats)	90 to	100	60 lbs.
Cow Peas (broadcast)	90 to	120	60 lbs.
Cow Peas (drilled)	50 to	60	60 lbs.
Soy Beans (broadcast)	60 to	120	60 lbs.
Soy Beans (drilled)	30 to	40	60 lbs.
Hairy Vetch (with Rye)	25 to	50	60 lbs.
Dwarf Essex Rape	4 to	7	50 lbs.
German Millet	40 to	50	50 lbs.
Hungarian Millet	40 to	50	50 lbs.
Japan Clover	5 to	6	25 lbs.
Sorghum	70 to	90	50 lbs.
Buckwheat	60 to	75	50 lbs.
Oats	75 to	90	32 lbs.
Barley	90 to	110	48 lbs.
Wheat	90 to	110	60 lbs.
Field Corn	8 to	10	56 lbs.
Rye	80 to	100	56 lbs.

Scott's Bacteria

BEFORE you can become enthusiastic about the inoculation of legumes you must know just what is to be gained by it. There is no better way to get a thorough understanding of the subject than to read our booklet "Friendly Workers of the Soil." It tells the whole story in 10 lessons clearly and interestingly. 8000 Schools of Agriculture used this booklet in the class-room last year. Instructors are pleased with it because of the clear and concise manner in which the subject is presented.



Our Bacteria chart of Questions and Answers will also give you some of the essential points. Those who do not believe in inoculation are, as a rule, not thoroughly informed about it. Those who know — inoculate. You owe it to yourself to read our booklet and chart. If you haven't copies, a post card is all you need to bring them.

Legume Bacteria

Ask for Bacteria "Questions and Answers"

NITROGEN, which is essential to the growth of all plants and animals, is constantly being removed from the soil. Some of the instrumentalities of its removal are: The growth of grain and other crops, the drainage of land, the action of wind and rain. A portion of this loss may be made up by the manure produced on the farm and by commercial nitrogen, but the cost of the latter is too great for profitable use. The only way in which nitrogen can be supplied so that farming may be profitably conducted is to draw upon the unlimited supply in the air.

Only the legumes, clover, beans, peas, etc., are capable of utilizing the nitrogen of the atmosphere. They are rich in protein, requiring more nitrogen than other plants; being heavy nitrogen feeders would be against them if they could not take the nitrogen from the air and use it. The air is made up of several gases, the proportion of nitrogen being 79%.

A legume cannot of itself make use of this nitrogen but the bacteria which live within the nodules on the roots are able directly to utilize the nitrogen of the air. The nitrogen passes into the bacteria and is changed along with other substances that are present into more complex nitrogenous substances which are used in some unknown way by the legume. The legume gives the bacteria a favorable place for development. It supplies the bacteria with sugars and other substances they need and in return the bacteria make the nitrogen of the air available for use of the plant.

The bacteria penetrate the root hairs and rapidly reproduce themselves. After a time the bacteria reach the interior of the root, still reproducing, and pass from one cell to another. The root enlarges and the nodule is formed. A plant cannot take nitrogen from the air unless the proper strain of bacteria is already in the soil or is introduced by inoculation.

To show that inoculation pays and performs a real service for the farmer legumes have often been grown experi-

mentally in sterilized soil that is entirely lacking in nitrogen. In these tests the inoculated plants make vigorous growth in spite of the absence of plant food in the soil.

Besides storing up nitrogen for the use of following crops, inoculation in most cases increases the growth of the inoculated legume, besides making this legume richer in protein.

The problem of keeping the soil as productive as possible is vital. If the ground is kept in a healthy condition you avoid half the gamble of farming. Inoculation is a big factor in increasing yields and in enhancing the value of your farm by building up the soil. The scientific way as compared with the soil method is not only more thorough but more economical.

"FRIENDLY WORKERS OF THE SOIL"

The subject of soil bacteriology and its relation to practical farming is interesting and important. There is too much to be said on this subject to do it justice in a few short paragraphs, so we have prepared a booklet entitled, "Friendly Workers of the Soil." You will not find it tedious reading and the information it contains should prove helpful to every farmer.

SCOTT'S GUARANTEED BACTERIA

The great advantages of Scott's Bacteria may be enumerated briefly: Ease of application, economy, and guaranteed results. These three factors are worthy of the closest consideration in buying inoculating material. It has been difficult in the past to secure cultures that measured up in all three respects.

In ordering please specify the kind of legume you want to inoculate. The following groups are each inoculated by different strains of bacteria: Alfalfa, Sweet and Bur Clovers; all true clovers, Red, Alsike, Mammoth, Crimson, and White; Cowpeas, Jap Clover, Lima and Velvet Beans; Soybeans; Garden and Field Beans; Garden and Field Peas, and Vetches.

If you have not received, "Friendly Workers of the Soil," or, having looked it over, wish additional copies, let us hear from you.

Prices

Scott's Bacteria is put up in cans. The smallest contain live Bacteria for the inoculation of thirty pounds of seed (price, 50c per can, postage paid); the standard size is for 60 pounds of seed, (price, one dollar postage paid); a special can brought out last year for Soybean cultures *only* is sufficient for five bushels of beans and sells at \$3.50 postage paid.

CONVENIENT ORDER BLANK

Many of our customers send in early spring orders for seed and bacteria. Later in the season when another legume, Soybeans, for example, is sowed, they forget about inoculation until the last minute and then rather than delay planting, go ahead and trust to good luck.

RUSH ORDER FOR SCOTT'S BACTERIA

Gentlemen:—

Please send immediately by prepaid parcel post the following order of Scott's Guaranteed Bacteria:

----- Cans for Sweet Clover or Alfalfa
----- Cans for Red, Alsike, Mammoth or Crim-
son.
----- Cans for Soybeans
----- Cans for Cow Peas or Jap Clover
----- Cans for Vetch or Field and Garden Peas
----- Cans for -----
Total ----- Cans.

Enclosed find check for \$----- Please send
cultures to

Name -----

Route -----

Post Office ----- State -----

SCOTT'S SEED GOES FARTHER

A FEW POINTS ABOUT ORDERING

Order Early. It always pays. Prices may be no higher, but sometimes it is impossible to get pure seed late in the season.

Order Blank. Use it, please. When shipping point is different from your mail address give county under each name.

WHERE IS MARYSVILLE

Marysville isn't a metropolis, so we often have to explain just where "The Town for Pure Seed" is located. Marysville is the County Seat of Union County, thirty miles from Columbus, on the C. C. C. & St. L. Railroad, between Springfield and Delaware and on the New York Central between Toledo and Columbus. We are in a better position than ever before to deliver seed quickly.

Parcel Post. Seed may be sent by parcel post according to the following table. In the first, second and third zone the weight limit is 70 pounds; in the others the weight limit is 50 pounds.

Owing to the new postal regulations it is necessary to add a 2-cent service charge on each package sent by parcel post. Please remember to add 2 cents for each package your order requires.

Zone	ZONE RATES	
	1st lb. or fraction	Additional lb. or fraction
1st within 50 miles of Marysville.....	7c	1c
2nd within 50 to 150 miles of Marysville.....	7c	1c
3rd within 150 to 300 miles of Marysville.....	8c	2c
4th within 300 to 600 miles of Marysville.....	9c	4c
5th within 600 to 1,000 miles of Marysville....	10c	6c
6th within 1,000 to 1,400 miles of Marysville..	11c	8c
7th within 1,400 to 1,800 miles of Marysville..	13c	10c
8th within 1,800 and over miles of Marysville..	14c	12c

BAGS

They are cheaper this year. If you allow for cotton bags in your order it will enable you to lay in a new supply at a reasonable price. We charge for bags what they cost us. We simply can't furnish bags free without adding their cost to the cost of the seed. We prefer to sell bags separately and make a separate item of them.

O. M. SCOTT & SONS COMPANY

Those who have a surplus of sacks may send their own and we will gladly use them. Be sure, however, that your name appears on the tags and that in the order you tell us the sacks are being sent.

CHANGE IN PRICE

All seed must be sold on the basis of market quotations. It is for this reason that our prices are for prompt acceptance. A slight fluctuation, however, does not affect our prices. We always accept orders at the prices quoted if at all possible, but we must follow any material changes whether they be up or down.

If you delay ordering after our price list has reached you some instructions for a possible adjustment should be sent. The simplest method is by adding or deducting seed. We often do the former or refund by check, but we never deduct seed unless so instructed. Taking the time to write causes a delay but we must be sure that the extra cost is approved by our customer.

Please remember that anything which detracts from efficiency is an expense and that eventually the buyer must pay it. We therefore ask your cooperation in enabling us to handle orders as economically and as promptly as possible. If you wish to do so just send along for the seed you require without writing for later quotations. Give us instructions to be followed in case of a change in price and we will guarantee that our treatment of the order will meet entirely with your approval.

CASH WITH ORDER

In Every Case We Must Adhere to This Policy

We would be perfectly willing to let down a little, but having had so much difficulty, especially in getting the small balances off our books, we are going to ask everybody, without exception, to include cash with order. If an error has been made in addition or if for any reason there is a small balance due, we will deduct seed. If you are not sure of price and in a hurry, you might send us a blank check, protecting yourself by writing in "not good for over \$-----," and we will fill in the cost of your order.

SCOTT'S SEED MAKES FEWER WEEDS

EMERGENCY ORDER BLANK

NAME _____

POST OFFICE.....

COUNTY _____ STATE _____

SHIPPING
POINT..... COUNTY.....

(If Different from P. O.)

SHIP BY: ☐ FREIGHT ☐ EXPRESS ☐ MAIL

We Ship by Freight Unless Otherwise Specified.

GENTLEMEN: DATE_____

Enclosed find \$_____for which ship the following:

BU.	LBS.	KIND OF SEED	Total lbs.	PRICE		TOTAL	
		Cans Bacteria for					
		Cans Bacteria for					
		BAGS					

TOTAL AMOUNT, \$-----

If you don't have recent prices when you want to make use of this blank, just send a check for what you believe will cover the order and the adjustment can be made by refund or invoice. You may rest assured the price will be consistent with the market. We'll guarantee it to be satisfactory.

PASS A GOOD THING ALONG

Every person who reads this Seed Guide has friends who would derive both pleasure and profit from it. Why not take it upon yourself to send a few names? That one section on "How To Know Good Seed," may help to cut down the annual crop of noxious weeds in the neighborhood.

Names.	Addresses.	Acres.

Sent by.....
Name.

We Pay the Freight

Last year for the first time we inaugurated the policy of paying freight on all orders of three hundred pounds or more to thirteen states. We are going to continue that plan during 1927. The states to which it applies are as follows: Ohio, Pennsylvania, West Virginia, New York, Illinois, Virginia, Michigan, Indiana, Kentucky, District of Columbia, Maryland, Delaware and New Jersey.

Outside those states we will allow 35 cents per one hundred pounds toward the transportation charges.

Please bear in mind that in spite of our reasonable quotations and the freight paid provision we commend Scott's Seed to you primarily on the grounds of superior quality. There is your real opportunity to save.

O. M. SCOTT & SONS COMPANY

Freight and Express Rates from Marysville, Ohio

	Freight Class		Ex. per 100 lbs.		Freight Class		Ex. per 100 lbs.
	3rd	4th			3rd	4th	
Arkansas				New York-Con.			
Little Rock.....	\$1.95	\$1.60	\$3.06	Delhi.....	\$0.70	\$0.48	\$2.03
California				Elmira.....	.61	.43	1.82
San Francisco.....	3.90	3.30	10.08	New York.....	.76	.53	2.23
Connecticut				Rochester.....	.56	.41	1.82
Hartford.....	.80	.57	2.39	Syracuse.....	.61	.43	1.92
Delaware				North Carolina			
Wilmington.....	.70	.48	2.13	Raleigh.....	1.28	.95	2.86
Dist. of Col.				Ohio			
Washington.....	.72	.50	2.03	Cambridge.....	.35	.27	1.09
Georgia				Celina.....	.34	.25	.94
Atlanta.....	1.68	1.52	2.75	Cincinnati.....	.39	.30	1.09
Illinois				Circleville.....	.31	.25	.94
Chicago.....	.55	.41	1.46	Cleveland.....	.44	.31	1.09
Danville.....	.52	.40	1.35	Mansfield.....	.34	.25	.94
Springfield.....	.60	.45	1.71	Pomeroy.....	.43	.33	1.09
Indiana				Portsmouth.....	.45	.34	1.46
Evansville.....	.52	.40	1.71	Toledo.....	.38	.29	.94
Indianapolis.....	.46	.35	1.35	Zanesville.....	.35	.27	1.19
South Bend.....	.53	.40	1.35	Pennsylvania			
Iowa				Clearfield.....	.72	.50	1.67
Des Moines.....	1.24	.97	2.50	Meadville.....	.70	.48	1.35
Kentucky				Philadelphia.....	.70	.48	2.13
Hickman.....	.71	.62	2.08	Pittsburgh.....	.51	.38	1.46
Lexington.....	.75	.63	1.35	Wellsboro.....	.72	.50	1.82
Williamsburg.....	.70	.48	1.97	Wilkesbarre.....	.70	.48	2.03
Maine				Rhode Island			
Portland.....	.80	.57	2.54	Providence.....	.80	.57	2.50
Maryland				South Dakota			
Baltimore.....	.73	.50	2.03	Pierre.....	1.54	1.18	3.85
Massachusetts				Tennessee			
Boston.....	.81	.57	2.39	Knoxville.....	.88	.70	2.39
Michigan				Vermont			
Detroit.....	.45	.34	1.19	Montpelier.....	.72	.50	2.39
Grand Rapids.....	.52	.40	1.35	Virginia			
Jackson.....	.46	.35	1.19	Charlottesville.....	.73	.51	2.03
Ludington.....	.70	.52	1.82	Richmond.....	.73	.51	2.39
Minnesota				West Virginia			
Minneapolis.....	1.07	.78	2.86	Charleston.....	.50	.37	1.61
Missouri				Clarksburg.....	.54	.40	1.46
Jefferson City.....	1.06	.81	2.44	Harrisville.....	.54	.40	1.19
New Hampshire				Huntington.....	.44	.33	1.46
Concord.....	.80	.57	2.54	Morgantown.....	.54	.40	1.61
New York				Wheeling.....	.51	.39	1.19
Albany.....	.73	.51	2.13	Wisconsin			
Buffalo.....	.56	.42	1.71	Madison.....	.61	.46	1.92
Canton.....	.80	.57	2.13	Milwaukee.....	.61	.46	1.82

